

5-1-2012

abd-A Regulation by the iab-8 Noncoding RNA

Maheshwar Gummalla
University of Geneva

Robert K. Maeda
University of Geneva

Javier J. Castro Alvarez
University of Geneva

Henrik Gyurkovics
Biological Research Center, Hungary

Kevin A. Edwards
Illinois State University

See next page for additional authors

Follow this and additional works at: <http://ir.library.illinoisstate.edu/fpbiosci>

 Part of the [Genetics Commons](#)

Recommended Citation

Gummalla, Maheshwar; Maeda, Robert K.; Alvarez, Javier J. Castro; Gyurkovics, Henrik; Edwards, Kevin A.; Singari, Swetha; and Bender, Welcome, "abd-A Regulation by the iab-8 Noncoding RNA" (2012). *Faculty Publications – Biological Sciences*. Paper 1.
<http://ir.library.illinoisstate.edu/fpbiosci/1>

This Article is brought to you for free and open access by the Biological Sciences at ISU ReD: Research and eData. It has been accepted for inclusion in Faculty Publications – Biological Sciences by an authorized administrator of ISU ReD: Research and eData. For more information, please contact ISURed@ilstu.edu.

Authors

Maheshwar Gummalla, Robert K. Maeda, Javier J. Castro Alvarez, Henrik Gyurkovics, Kevin A. Edwards, Swetha Singari, and Welcome Bender

abd-A Regulation by the *iab-8* Noncoding RNA

Maheshwar Gummalla¹, Robert K. Maeda¹, Javier J. Castro Alvarez¹, Henrik Gyurkovics², Swetha Singari³, Kevin A. Edwards³, François Karch^{1*}, Welcome Bender^{4*}

1 Department of Genetics and Evolution, University of Geneva, Geneva, Switzerland, **2** Institute of Genetics, Biological Research Center, Szeged, Hungary, **3** School of Biological Sciences, Illinois State University, Normal, Illinois, United States of America, **4** Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts, United States of America

Abstract

The homeotic genes in *Drosophila melanogaster* are aligned on the chromosome in the order of the body segments that they affect. The genes affecting the more posterior segments repress the more anterior genes. This posterior dominance rule must be qualified in the case of *abdominal-A* (*abd-A*) repression by *Abdominal-B* (*Abd-B*). Animals lacking *Abd-B* show ectopic expression of *abd-A* in the epidermis of the eighth abdominal segment, but not in the central nervous system. Repression in these neuronal cells is accomplished by a 92 kb noncoding RNA. This “*iab-8* RNA” produces a micro RNA to repress *abd-A*, but also has a second, redundant repression mechanism that acts only “*in cis*.” Transcriptional interference with the *abd-A* promoter is the most likely mechanism.

Citation: Gummalla M, Maeda RK, Castro Alvarez JJ, Gyurkovics H, Singari S, et al. (2012) *abd-A* Regulation by the *iab-8* Noncoding RNA. *PLoS Genet* 8(5): e1002720. doi:10.1371/journal.pgen.1002720

Editor: Norbert Perrimon, Harvard Medical School, Howard Hughes Medical Institute, United States of America

Received: November 14, 2011; **Accepted:** April 3, 2012; **Published:** May 24, 2012

Copyright: © 2012 Gummalla et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a grant from Otko to HG; a grant from the National Institutes of Health to KAE; grants from the Swiss National Fund for Research, the NCCR Frontiers in Genetics, the State of Geneva, and the Foundation Claraz to FK; and a grant from the National Institutes of Health to WB. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Francois.Karch@unige.ch (FK); heber_domingues@hms.harvard.edu (WB)

Introduction

Genome wide surveys for RNA transcription units in a variety of eukaryotes have revealed a surprising number of transcripts that are not traditional messenger RNAs. A variety of functions have been suggested for these “noncoding” RNAs (ncRNAs), although the large majority have no known purpose (reviewed in ref [1]). In *Drosophila melanogaster*, primary transcripts cover at least 60% of the genome [2]. Many of these transcripts do not correspond to defined genes, but they are evolutionarily conserved. Particular attention has been given to ncRNAs in the bithorax complex (BX-C), a cluster of three homeobox-containing transcription factors required for segment identity (reviewed in ref. [3]). Although much of the ~300 kb of BX-C DNA is transcribed, the BX-C contains only one other protein coding sequence [4]. Lipshitz et al. [5] first described apparent ncRNAs from the BX-C in the *bithoraxoid* regulatory region. They suggested that such transcripts could reflect nonspecific initiation by RNA polymerase near a strong enhancer, a possibility that still remains attractive. Several other ncRNAs in the BX-C have been identified by Northern blots or RNA in-situ hybridizations [6–8]. It has been suggested that such transcripts might block silencing by the Polycomb Group proteins [9], but this idea is not yet supported by the analysis of existing mutations. A readthrough product of the *bithoraxoid* ncRNA transcription unit may repress features of early transcription from the *Ultrabithorax* (*Ubx*) promoter [10], and the *iab-4* and *iab-8* ncRNAs are the likely precursors for micro RNAs (miRNAs) [11–13]. Otherwise, these ncRNAs still lack functions.

Prior indications of the *iab-8* noncoding RNA

Here, we focus on the *iab-8* ncRNA. Several lines of evidence have suggested the existence of a 90 kb-long transcription unit, extending between *Abd-B* and *abd-A*, with a likely start site within the *iab-8* regulatory region. RNA in-situ hybridizations to embryos, using genomic DNA fragment from the *iab-2* through the *iab-8* regulatory regions as probes, detect an RNA in the 8th and 9th abdominal segments (parasegments 13 & 14). Strand-specific probes revealed that it is transcribed in a distal-to-proximal direction (from *Abd-B* towards *abd-A*) [7,8,14,15]. This transcript is first seen at about stage 6 [11] in the epidermis, but from stage 14 onward (germband shortening), the RNA is detected only in the developing central nervous system (CNS). A promoter for an uncharacterized RNA was independently mapped to the *iab-8* region, just downstream of the *Abd-B* transcription unit [16].

Additionally, a transcript starting in the *iab-8* region has been suggested as the precursor for a micro RNA, called miR-*iab-8* or miR-*iab-4AS* [11–13]. This miRNA is transcribed from the *iab-3* regulatory region in the distal-to-proximal direction, and strand specific genomic probes from this region indicate that the precursor is made in the 8th and 9th abdominal segments, as described above. This miRNA is required for male and female fertility, and complementation tests with a series of rearrangement breakpoints suggest that the start site of this RNA is in the *iab-8* regulatory region, downstream of *Abd-B* [11].

Here, we characterize the structure and function of the 92 kb long “*iab-8* ncRNA”. This ncRNA represses the expression of the homeotic gene *abd-A* in the posterior CNS. This repression

Author Summary

Although long, noncoding RNAs have been found in many organisms, it has been difficult to assign to them any molecular function. The homeotic gene clusters in the fruit fly, *Drosophila melanogaster*, contain many such noncoding RNAs. We have characterized one such noncoding RNA, a 92 kb transcription unit from within the bithorax complex. This transcript, called the *iab-8* ncRNA, is made in the cells of the central nervous system in the eighth abdominal segment, along with the homeotic transcription factor *Abdominal-B*. Another homeotic transcription factor, *abdominal-A*, is repressed in these cells. It has generally been assumed that *abdominal-A* repression in these cells is mediated by the *Abdominal-B* protein. However, here we show that it is not *Abdominal-B* that represses *abdominal-A*, but the *iab-8* ncRNA. This repression is accomplished by two redundant mechanisms; the *iab-8* precursor produces a micro RNA, which targets the *abdominal-A* mRNA, and *iab-8* transcription interferes with the *abdominal-A* promoter, which lies just downstream of the *iab-8* ncRNA poly(A) site.

depends not only on the miR-*iab-8* micro RNA, but also on transcriptional interference in the region of the *abd-A* promoter.

Results/Discussion

Repression of *abd-A* in the 8th abdominal segment

In wild type embryos, *abd-A* expression is detected in the epidermis and CNS of PS7 to PS12 but not in PS13 (Figure 1A). *Abd-B* is strongly expressed in PS13, and it was initially claimed that *Abd-B* represses *abd-A* in PS13 [17], just as *abd-A* represses *Ubx* and *Ubx* represses *Antp* [18]. This repression hierarchy can account for the dominance of posterior homeotic genes over anterior ones, often called “posterior prevalence” [19]. Indeed, embryos homozygous for *Df(3R)C4*, which removes *Abd-B*, show ABD-A expression extending throughout PS13 (Figure 1B). However, the *Df(3R)C4* deficiency extends downstream of the ABD-B transcription unit, removing all of the *iab-8* regulatory region and part of *iab-7* (Figure 2). Surprisingly, embryos homozygous for an *Abd-B* null point mutation, *Abd-B^{D16}*, show ABD-A derepression in PS13 of the epidermis, but not in the CNS (Figure 1C). Homozygotes for *Abd-B^{D18}*, a deletion removing all of the *Abd-B* coding sequences (Figure 2), show the same ABD-A expression pattern (not shown). This unexpected repression of ABD-A in the CNS can be seen most dramatically in the *Abd-B^{D14}* mutation. *Abd-B^{D14}* deletes the promoter for the *Abd-B* “m” transcript [20], expressed from PS10 through PS13, but leaves the promoters for the “r” transcripts expressed in PS14. In the CNS of *Abd-B^{D14}* homozygotes, *abd-A* does not fill in the gap left by the absence of *Abd-B* in PS13 (Figure 1D). Clearly then, there must be some function deleted by *Df(3R)C4* that is not affected by *Abd-B^{D18}* or more subtle *Abd-B* mutations. Our attention turned to the *iab-8* ncRNA, which appeared to initiate in the *iab-8* region deleted in *Df(3R)C4*.

Mapping the *iab-8* ncRNA exons

The spliced product of the *iab-8* ncRNA was initially uncovered by a fortuitous insertion of an exon-trap mobile element. This element, a derivative of the *Mimos* mobile element, is called Hostile takeover (*Mi(Hto-WP)*; Genbank #JN049642). An insertion was recovered in the *iab-6* domain of the BX-C (“TA” target site bases 85,277 & 85,278), named *Mi(Hto-WP)LNP* or simply *LNP*, for short (Figure 2). 3' RACE products were amplified with primers




Figure 1. ABD-A expression in *Abd-B* mutant embryos. Stage 14 embryos in A–C were stained with antibody to ABD-A, opened along the dorsal midline and flattened for photography. ABD-A is absent from PS13 in wild type (A), but appears throughout PS13 in *Df(3R)C4* homozygotes (B). In *Abd-B^{D16}* homozygotes, ABD-A is only in the lateral and dorsal epidermis of PS13 (C). Dissected CNS's from stage15 embryos in D were doubly stained for ABD-A (red) and ABD-B (green). In wild type, the expression domains overlap through PS10–12, with some nuclei expressing both proteins. In *Abd-B^{D14}* homozygotes, ABD-B expression is absent from PS10–13, but the ABD-A pattern is unchanged, leaving PS13 without either protein.
doi:10.1371/journal.pgen.1002720.g001

within *LNP* and within the 3' exon of *abd-A*. The sequence of the product revealed the exon structure diagrammed in Figure 2. The sequence included 5 novel exons before it spliced into *abd-A*, at the 5th exon of the predominant splice form of the *abd-A* mRNA [4,17]. Many of these exons match those of a cDNA designated MIP06894 (Genbank BT099824.1)(Figure 2), identified by the Berkeley *Drosophila* Genome Project.

The exons of the *LNP* cDNA were used to generate primers for 5' and 3' RACE, using total RNA from Oregon R embryos. Figure 2 diagrams the predominant splicing product, which spans ~92 kb. An RT/PCR product was recovered and sequenced that extended from exon 1 through exon 8, as well as one that extended from exon 1 through exon 7, and then included exons 5, 6, and 7 of the *abd-A* transcription unit. Figure 2 also shows three alternate 5' exons and five alternate 3' splicing patterns. RT/PCR products included extensions of exons #2 or 4, ending at sites of genomic poly(dA) stretches; these were likely derived from splicing intermediates. Rare clones were also recovered that skipped exons, splicing from exons 1, 2, or 6 into *abd-A* exons 5 or 6. Exon 1 had two start sites separated by 135 bases; the upstream start was ~3 fold more abundant. Exon 4 included only 6 bases, although rare products included an alternate 3' extension of 92 bases. Quantitative PCR was also used to show that termination at exon 8 was ~500-fold more common than splicing into *abd-A*. The sequences of the predominant and alternate exons are given in Figure S1. Two recent genome-wide searches for novel noncoding transcripts in embryos have uncovered some of these same




Figure 2. Map of the abdominal half of the bithorax complex. The horizontal bar indicates the DNA sequence map, numbered in kb according to Martin et al. [4] (Genbank U31961). Base #1 corresponds to base 12,809,162 on chromosome 3R in release 5.37 of the *Drosophila* genome. The coordinates proceed distal to proximal on chromosome 3R, which is opposite in orientation to the whole genome numbering. The regulatory domains *iab-2* through *iab-8* are color coded; the domain borders are defined by deletion mutations (*Fab8* [22]; *Fab7*, [41]; *Mcp*, [44]; *iab-3/iab-4*, L. Sipos personal communication), or inferred from the binding sites of the CTCF factor [45]. Below the DNA bar are shown the splicing patterns of *abd-A* and *Abd-B* (in black), a cDNA derived from the *Mi(Hto-WP)LNP* insertion (red), and the MIP06894 cDNA (green). At the bottom, the splicing pattern for the *iab-8* ncRNA is shown in dark blue, with numbered exons, and alternate 5' or 3' extensions indicated with light blue lines. Mutant lesions are indicated above the DNA bar. The rearrangement breakpoints are color coded according to their phenotypes when heterozygous with the *mfs5649* insertion.
doi:10.1371/journal.pgen.1002720.g002

transcripts [2,21]. Graveley et al. [2] also reported transcripts from adult males with most of the same exons, but an alternate start site, in the *iab-6* region.

The promoter for the *iab-8* RNA maps distal to the *Fab-8* boundary, in the *iab-8* regulatory region [22]. The *iab-8* region should be under *Polycomb* Group repression in parasegments 1–12, which explains why the transcript is only expressed in PS13 and 14 [11]. Exons 1–7 appear to be evenly spaced across the abdominal region of the bithorax complex, with one in each of the *iab* cis-regulatory domains. A comparison with the genomic sequences of various *Drosophila* species suggests that the sequences of the exons are not more conserved than those of the introns. However, the existence of the exons does appear to be conserved, in that the splice junctions are among the most conserved features of the exons. This is illustrated in Figure 3A for exon 3, in the *iab-6* region. The embryonic expression pattern is also conserved;

expression is restricted to PS13 and 14 in *D. pseudoobscura* and *D. virilis*, as it is in *D. melanogaster* (see Figure S2).

The spliced product of the *iab-8* RNA is non-coding by traditional criteria, but the possibility of small peptides [23,24] cannot be ruled out. In particular, exon 8 includes a potential 9 amino acid peptide, with appropriate translation initiation and termination signals, and the coding potential for this peptide is well conserved in *D. ananassae*, *D. pseudoobscura* and *D. willistoni* (Figure 3B), although it is not found in *D. virilis* and more distantly related species.

abd-A repression by the iab-8 RNA

There are many chromosome rearrangements, mostly from the collection of E. B. Lewis, which interrupt the *iab-8* ncRNA transcription unit. These can be used to test whether truncated versions of the *iab-8* RNA can repress *abd-A*. Rearrangement

A.

Exon 3

<i>D. melanogaster</i>	TATGCCCTTCCAGTTTGATTACACATCGACCCCTGGAGCGAGGACTTAACCCGACGGCAGCTGCATCAGTTGAGTCTCCATCTTAACATA
<i>D. erecta</i>	TAGCCATTCCAGTTTGATTACACATgGACCCCTGGAGtGAGGACTTAACCCGACGGCAGCTGCATCAGTTGAGTCTCCATCTcAgCgct
<i>D. ananassae</i>	TtaaccTTCCAGTTTGATTAtcCaacGACCCCTGGAGtGAG-----CGcCGGCAGCTGCATCAGTTGAGT-----
<i>D. willistoni</i>	TtTgtCaTTtCAGTTgGATT-----gCtatTGGg-----aGCAGCTGttacAGTtAGT-TcctTccAAt----

B.

	M	L	N	L	Q	F	M	S	N	*	*		
<i>D. melanogaster</i>	CAA	ATG	CTT	AAT	TTA	GAA	TTT	ATG	AGC	AAT	TAA	TTT	TAA
<i>D. ananassae</i>	CAA	ATG	TTT	AAT	TTG	GAA	TTT	ATG	AGT	AAT	TAA	TTT	TAA
			F		L				S				

Figure 3. Evolutionary conservation. A. A comparison of exon 3 and neighboring bases with the homologous regions from the genomes of three other *Drosophila* species. B. Potential nine amino acid peptide within exon 8 of the *iab-8* ncRNA. The *D. melanogaster* sequence is compared to that of *D. ananassae*. The initial methionine codon is preceded by a perfect translation start consensus sequence [46], and there are two stop codons after the 9th amino acid. The three bases altered in *D. ananassae* are highlighted in red; only one changes the predicted amino acid.
doi:10.1371/journal.pgen.1002720.g003

breaks that truncate the *iab-8* RNA near its start site cause a dramatic derepression of *abd-A* in the CNS of the 8th abdominal segment, indistinguishable from that seen in *Df(3R)C4* homozygotes (Figure 1). Rearrangements with this effect include *iab-7^{SGA62}*, *iab-6¹¹*, *iab-5^{B43}*, *iab-4²³³⁰*, and *iab-4¹⁸⁶* (Figure 2 & 4). The same spread of ABD-A into the CNS of PS13 is seen with the *Fab-8⁶⁴* deletion, which removes the *iab-8* ncRNA promoter (Figure 2). Interestingly, embryos homozygous for chromosome breaks mapping closer to *abd-A* show a much more subtle derepression of *abd-A*. In *iab-3⁵⁰²²* homozygotes, for example, weak misexpression is limited to a few cells (Figure 4). Similar weak misexpression is seen in homozygotes of *iab-4,5³²⁴⁵* and *iab-4³⁰²* (Figure 2). Finally, embryos homozygous for the *iab-3³³* rearrangement show *abd-A* misexpression in only a very few CNS cells in the most anterior part of the 8th abdominal segment (Figure 4). This break lies downstream of the poly(A) addition site of the major *iab-8* transcript, but upstream of the *abd-A* transcription start site.

The difference between the two classes of breakpoints seems to be the expression of miR-*iab-8*. The *iab-4¹⁸⁶* break, maps just upstream (within 3 kb) of the miR-*iab-8* coding region and shows complete loss of *abd-A* repression in PS13. In contrast, the *iab-3⁵⁰²²* break maps ~5 kb downstream of miR-*iab-8* and shows only slight misexpression. Thus, one might guess that miR-*iab-8* is responsible for most of the repression of *abd-A*, especially since the 3' UTR of *abd-A* includes sequences homologous to the "seed" region of miR-*iab-8* [12,13]. However, embryos homozygous for a deletion of miR-*iab-8* (Δ miR-*iab-8*) do not show a dramatic misexpression of ABD-A in the PS13 CNS [11]. A closer examination of these homozygous embryos does reveal a weak misexpression of *abd-A* in a small number of nuclei in anterior PS13 (Figure 4), but clearly not the strong and widespread misexpression of *iab-4¹⁸⁶*. Thus, it appears that miR-*iab-8* does repress *abd-A* in the PS13 CNS, but there must be a second, redundant function of the *iab-8* RNA to completely represses *abd-A*. UBX expression in embryos is apparently not affected by this

second function; its expression pattern in *iab-7^{SGA62}* homozygous embryos is the same as that in miR-*iab-8* deletion homozygotes (not shown).

Fertility function of the *iab-8* ncRNA

A deletion of the miR-*iab-8* causes sterility in both sexes [11]. Thus, we expected that any combination of alleles that failed to make the miR-*iab-8* micro RNA would be sterile, including, for example, an *iab-7* break (*iab-7^{MX2}* or *iab-7^{SGA62}*) heterozygous with Δ miR-*iab-8* [11]. There is an insertion of the "PZ" P element ~4.2 kb downstream of the *iab-8* RNA start site, designated *mfs(3)05649* (here called *mfs5649*; Figure 2). Homozygotes are sterile in both sexes, and the females show the same phenotype (blockage of the oviduct) as is seen in Δ miR-*iab-8* homozygotes [25]. We assume the *mfs5649* insertion truncates the *iab-8* RNA, since it fails to complement with Δ miR-*iab-8* for the sterility phenotype. The *Fab8⁶⁴* deletion (derived from the *mfs5649* P element; Figure 2; [22]) is also sterile as a homozygote or as a heterozygote with Δ miR-*iab-8*.

We tested rearrangement breakpoints in the *iab-2,3*, and *4* regions, downstream of the miR-*iab-8* template, for fertility when heterozygous with the *mfs5649* P element. Surprisingly, many rearrangement breakpoints 3' to the miR *iab-8* template have a female sterility phenotype when heterozygous with *mfs5649* (Figure 2); males of these genotypes are fertile. These sterile females show a failure of mature oocytes to move through the oviduct, much like *mfs5649* homozygotes or the Δ miR-*iab-8* homozygotes. It does not seem likely that breakpoints downstream of the miR-*iab-8* template interfere with the proper processing of the micro RNA, because these same breakpoints are fertile when heterozygous to Δ miR-*iab-8*. It is possible that the subtle misexpression of ABD-A in PS13 seen in *iab-3* breaks is responsible for the female sterility, especially if the misexpression it is more dramatic at later times in development. Not all breakpoints give this female sterility phenotype, and there is no apparent order to




Figure 4. ABD-A expression in rearrangements truncating the *iab-8* ncRNA. Embryos homozygous for the indicated mutations were doubly stained for ENGRAILED (green) and ABD-A (red), and the CNS's were dissected and photographed. The posterior end of each CNS is shown; the ENGRAILED stripes mark the parasegmental boundaries. The *iab-4* and *iab-7* breaks cause widespread misexpression of ABD-A in PS13, but *iab-3* breaks show only subtle misexpression in a few nuclei. Embryos homozygous for a deletion of the *iab-8* miRNA also show misexpression in only a few nuclei.

doi:10.1371/journal.pgen.1002720.g004

the fertile and sterile breakpoint alleles (Figure 2). Some of the rearrangements may fuse the *iab-3* region with novel transcription units, restoring the repression of *abd-A* in the critical cells.

Mechanism of repression

The *iab-8* ncRNA could make a product, such as another miRNA, that represses *abd-A*. Indeed, there is a secondary structure hairpin in exon 6 of the spliced transcript that could serve as a miRNA precursor. The *iab-8* ncRNA could also code for tiny peptides, as noted above (Figure 3B). These possibilities prompted us to misexpress the *iab-8* ncRNA spliced product. A cDNA cassette, representing the major splicing product (Figure 2) plus 236 bp of genomic DNA downstream of the poly(A) addition site, was cloned into the pUAST vector [26]. P element transgenes were recovered and crossed to flies expressing the yeast GAL4 activator in abdominal segments 3–8 (parasegments 8–13). However, embryos containing both the GAL4 activator and the UAS/*iab-8* cDNA target showed no apparent reduction in the ABD-A levels in the segments expressing GAL4 (not shown).

The cDNA misexpression experiment does not rule out a product made from an intron, such as an RNA component of a diffusible repressive complex, as alleged for non-coding RNAs in mammalian HOX complexes [27]. If the putative second repressor involves a diffusible molecule, it should be able to act on both chromosomes, even if it is only produced by one. The miR-*iab-8* micro RNA should be diffusible in this way, and so, to examine the second repressive function, we needed to test genotypes lacking miR-*iab-8*. Specifically, heterozygotes were made with the Δ miR-*iab-8* deletion on one chromosome, and with a mutation truncating the *iab-8* RNA upstream of the miRNA template (*mfs5649*, *iab-7^{SCA62}*, or *iab-5^{B45}*) on the other chromosome. Such embryos make the *iab-8* RNA from only one chromosome, and cannot make the micro RNA from either. As shown in Figure 5, these embryos showed strong ABD-A




Figure 5. Test for *trans* repression by the *iab-8* ncRNA. CNSs, stained for ABD-A, were dissected from embryos of the indicated genotypes. In Δ miR/*mfs5649* embryos, only one chromosome makes a full length *iab-8* ncRNA, and neither chromosome produces the *iab-8* miRNA, as diagrammed. The strong expression of ABD-A in PS13 in this genotype shows that the *abd-A* gene on the *mfs5649* chromosome is not repressed, i.e. the *iab-8* ncRNA acts only in *cis*. doi:10.1371/journal.pgen.1002720.g005

misexpression in the CNS of PS13 (the 8th abdominal segment), suggesting that the *iab-8* RNA can only repress the copy of *abd-A* on the chromosome from which it is transcribed. To control for a potential effect of haploinsufficiency of the *iab-8* RNA, the Δ miR-*iab-8* deletion was also tested over *DjP9*, a deletion that removes the entire bithorax complex. These Δ miR-*iab-8*/*DjP9* embryos show no apparent misexpression of ABD-A in PS13. Thus, the second *iab-8* RNA repressive function must act only in *cis*.

In a similar test, we employed a duplication for the proximal two thirds of the complex, Dp(3:2)D109, which extends into the *iab-5* region (at \sim 110 kb; [28]). This duplication includes *abd-A*, but lacks the *iab-8* RNA promoter. Embryos homozygous for the Δ miR-*iab-8* deletion but containing this duplication also show ABD-A misexpression in the PS13 CNS (Figure S3). Thus, there are two mechanisms by which the *iab-8* RNA represses *abd-A*, first, through production of the *iab-8* miRNA (acting in *trans*), and second, a repressive function acting only in *cis*. The Supplementary Table S1 summarizes which genotypes supply which repressive functions.

The *cis*-repression of one transcription unit by another is often termed transcriptional interference. This term, however, encompasses several possible molecular mechanisms [29]. An example of a long, ncRNA involved in transcriptional *cis*-repression is the XIST RNA, involved in mammalian X chromosome inactivation [30] (A recent report suggests that the XIST RNA can also work in *trans* [31]). Nascent transcripts are involved in repression in RNAi silencing of heterochromatin in fission yeast [32] and in RNA-directed DNA methylation in *Arabidopsis* [33]. By analogy to these systems, the *iab-8* RNA could recruit gene silencing machinery to the site of its transcription. The RNA sequences required for such recruitment might be mapped by examination of deletions in the BX-C. Ideally, the *iab-8* miRNA should be removed to have a clear assay for the *cis* repressor. We have checked embryos homozygous for the *Fab3,5^{DV}* deletion (Figure 2), which covers the site of the *iab-8* miRNA precursor; they still show *abd-A* repression in the posterior CNS. Likewise, a double deletion chromosome, with Δ miR-*iab-8* and *Fab7^L*, also retains the *cis* repression. The *Fab7^L* deletion (Figure 2) was tested because it removes a Polycomb Response Element [34,35] which is coincident with exon 2 of the cDNA. Two other deletions have been examined which span the *iab-4* through *iab-7* regions, although both retain the *iab-8* miRNA (*iab-4,5,6^{DB}* and *iab-6,7^{HL}*; Figure 2). In these, we looked for more subtle misexpression, such as that seen in *iab-3* breaks (Figure 4), but no such misexpression was seen. This analysis does not yet cover the *iab-2* and *iab-3* regions, nor does it exclude the possibility of multiple redundant sequences throughout the transcription unit that could recruit repressive factors.

A more likely repression mechanism, perhaps, is that the RNA polymerase transcribing the *iab-8* RNA somehow interferes with the *abd-A* promoter. Examples of this type of transcriptional interference come from budding yeast, where the GAL7 gene is repressed by the upstream GAL10 transcript [36], and the SER3 gene is repressed by the upstream, noncoding SRG1 transcript [37]. In these cases, the 3' ends of the upstream transcripts are close to the downstream promoters, suggesting repression by occlusion of the downstream promoters or their proximal enhancers. If the *iab-8* RNA interferes with an *abd-A* enhancer, that enhancer must lie downstream of the *iab-4¹⁸⁶* breakpoint, since *abd-A* is totally derepressed in the PS13 CNS in embryos homozygous for this break (Figure 4). The *abd-A* promoter seems like the most likely target of interference, since the major poly(A) site of the *iab-8* RNA lies only 1.1 kb upstream of the initiation site of *abd-A*, and the *iab-8* RNA primary transcript likely continues past its poly(A) addition site [38]. In any case, minor splice variants

clearly do continue past the poly(A) site and into the *abd-A* transcription unit (Figure 2).

The function of the *iab-8* ncRNA fits with the rule of posterior dominance - it blocks expression of a more anterior homeotic gene in more posterior segments. The repression of *Ubx* by the *bxl* ncRNA [10], although subtle, fits the same pattern. The novel aspect, here, is that this posterior repression can be accomplished by noncoding transcription units, in addition to DNA binding proteins. The mechanism of transcriptional interference would fix the arrangement of these ncRNAs in the bithorax complex. It seems possible that the ancestral HOX complex turned off anterior genes by readthrough transcripts of more posterior genes, or by noncoding RNAs initiating from their posterior enhancers. Such a method of repression would dictate the linear order of the HOX genes, 3' to 5', anterior to posterior.

Materials and Methods

Drosophila strains

Wild type stocks were Canton *S* or Oregon R. Mutations included *abd-A*^{MX1}, *iab-3*²⁷⁷, *iab-4*³⁰², *iab-5*⁸⁴³, *iab-7*^{SGA62}, *iab-7*^{MX2}, *Abd-B*^{D16}, *Abd-B*^{D14}, *Df(3R)C4*, *Df(3)P9* (ref [39]); *iab-3*³³, *iab-3*^{B1} (ref [17]) *iab-6*¹¹ ref [40]; *Fab7*¹ ref [41]; *mfs(3)05649* (ref [25]), *Fab8*⁶⁴ ref [22]; *Fab3-5*^{DV} ref [42]; *iab-3*⁸⁶⁴, *iab-3*⁵⁰²², *iab-4*¹⁸⁶, *iab-4*²³³⁰, *iab4,5*³²⁴⁵, *iab-5*⁸⁴³, Δ *miR-iab-8* (ref [11]); *T(3:2)DpD109* [28] and *Mi[Hto-WP]* (described here).

Antibody staining

Embryos were fixed, stained, and mounted as described by [17]. Primary antibodies used were mouse anti-ADB-B (1:2 dilution, developed by S. Celniker, Developmental Studies Hybridoma Bank), mouse anti-UBX (1:10, developed by R. White, Developmental Studies Hybridoma Bank), rabbit anti- β -galactosidase (1:1500, Cappel/MP Biomedicals), mouse anti- β -galactosidase (1:1000, Promega), rabbit anti-En (1:500, Santa Cruz Biotechnology), mouse anti-ABD-A (1:500, 6A18.12, gift of I. Duncan), and goat anti-ABD-A (1:100, Santa Cruz Biotechnology). Secondary antibodies were donkey anti-mouse, donkey anti-goat, and donkey anti-rabbit, coupled to either Alexa 488 or Alexa 555 (1:500, Invitrogen), and HRP coupled goat anti-mouse (1:1000, Bio-Rad).

The CNS's were hand dissected with tungsten needles and placed on a glass slide in a drop of Immu-Mount (for HRP staining, Shandon) or Vectashield with DAPI (for fluorescence, Vector Laboratories), and then gently flattened under a coverslip. Fluorescence images were taken with a Leica SP2 AOBS confocal microscope; the fluorescence pictures show free projection averages of stacks of images, after scanning through the depth of the tissue. Homozygous embryos were identified by the absence of lacZ staining from the *TM3 ftz-LacZ* balancer.

Fertility tests

Each of ten mutant virgin females was placed in a vial with three wild type males. Likewise ten mutant males were mated, each with three wild type virgin females. Vials were maintained at 25° for five days, and then examined for the presence of larvae.

cDNA analysis

Adults heterozygous for *Mi[Hto-WP]* and *Hsp70-Gal4* (Bloomington stock #1799) were heat shocked for 45 min. at 37° to induce GAL4 expression, and then left to express the LNP transcript at room temperature for 4 h. RNA was then isolated using TRI reagent (Sigma) and reverse transcribed with MMLV reverse transcriptase (Promega) using an adaptor primer (GAA-

GACAGACACCGGACT18V). PCR was then performed using a forward primer in *Hto* and a reverse primer in the 6th exon of *abd-A*. The resulting amplicon was sequenced to identify the splicing pattern.

Total RNA from Oregon R embryos was prepared using the RNAAqueous-4PCR kit (Ambion), and 3' RACE and RNA ligase-mediated 5' RACE reactions were performed using the First-Choice RLM-RACE kit (Ambion). The 5' RACE procedure was designed to recover only capped 5' ends. Gel-isolated products were sequenced directly, or cloned first into the PCR-Blunt vector (Invitrogen). Quantitative PCR reactions used cDNA prepared from 6–12 h old embryos. The initial cDNA products were compared to measured dilutions of amplified cDNA products covering the relevant exons.

RNA in situ hybridization and embryo staining

The production of digoxigenin-labeled probes and the hybridization of embryos was as described by Fitzgerald and Bender [11], except that acetone treatment [43] was used instead of proteinase K for permeabilization of the embryos. Clones spanning exon 8 from *D. melanogaster*, *D. pseudobscura* and *D. virilis* were recovered after PCR reaction on genomic DNAs with the following pairs of oligonucleotides: *D. melanogaster* 5'CGCTCGA-GAGATTACAAACG3' and 5'GGTGTATTACGGTCAAG-GGGG3' generating a fragment of 1013 bp; *D. pseudobscura* 5'CAGGCATTCAGTAAACACGGC3' and 5'GGATGTGTC-GAGTGGTGTGG3' generating a fragment of 1477 bp; *D. virilis* 5'CTTTCGGTCTTATTCAACGG3' and 5'CCGATCCTGCTGGTGTGTC3' generating a fragment of 1364 bp.

Supporting Information

Figure S1 Sequences of *iab-8* ncRNA exons. The first and last bases of each exon are numbered according to the SEQ89E coordinates of Martin et al. [4] (Genbank U31961). (PDF)

Figure S2 Conserved *iab-8* noncoding RNA expression patterns in *D. melanogaster*, *D. pseudobscura* and *D. virilis* embryos. The top 3 panels show embryos at stage 8, while the bottom panels show embryos at stages 14–17. (TIF)

Figure S3 Additional test for *trans* repression by the *iab-8* ncRNA. Males of the genotype *T(2;3) DpD109, AmiR, Fab/TM2* were crossed to Δ *miR/TM3, ftz-LacZ* females. The *TM3*-containing embryos were recognized by their LacZ expression. Among the remaining embryos, half showed no apparent ABD-A expression in the CNS of PS13 (presumed to be *TM2/AmiR*), and half gave clear PS13 misexpression (presumed *DpD109, AmiR/AmiR*). Thus, the repression fails to act in *trans* on the duplication. The PS13 misexpression is weaker than the PS7-12 level, because the former derives from only the one *abd-A* copy on the duplication, but the latter represents three doses of the *abd-A* gene. *DpD109, +/AmiR* embryos produced in a control cross displayed little, if any, ABD-A misexpression in PS13. (TIF)

Table S1 Two mechanisms, *iab-8* miRNA *trans*-repression and *cis*-repression mediate *abd-A* repression in PS13 of the CNS. The table summarizes which of these 2 mechanisms is/are affected in the various mutant alleles. Note that complete ectopic expression in PS13 is only observed when both mechanisms are affected. (TIF)

Acknowledgments

We are grateful to Ian Duncan for the gift of anti-ABD-A and for the *DpD109* strain. We thank a reviewer for suggesting the *DpD109* experiment. We thank Eva Favre and Jorge Faustino for excellent technical assistance. The antibodies from the Developmental Studies Hybridoma Bank were developed under the auspices of the NICHD and maintained by the University of Iowa, Department of Biology.

References

- Ponting CP, Oliver PL, Reik W (2009) Evolution and functions of long noncoding RNAs. *Cell* 136: 629–641.
- Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, et al. (2011) The developmental transcriptome of *Drosophila melanogaster*. *Nature* 471: 473–479.
- Maeda RK, Karch F (2006) The ABC of the BX-C: the bithorax complex explained. *Development* 133: 1413–1422.
- Martin CH, Mayeda CA, Davis CA, Ericsson CL, Knafels JD, et al. (1995) Complete sequence of the bithorax complex of *Drosophila*. *Proc Natl Acad Sci U S A* 92: 8398–8402. Issn: 0027–8424.
- Lipshitz HD, Peattie DA, Hogness DS (1987) Novel transcripts from the Ultrabithorax domain of the bithorax complex. *Genes Dev* 1: 307–322.
- Cumberledge S, Zaratzian A, Sakonju S (1990) Characterization of two RNAs transcribed from the cis-regulatory region of the abd-A domain within the *Drosophila* bithorax complex. *Proc Natl Acad Sci U S A* 87: 3259–3263.
- Sanchez-Herrero E, Akam M (1989) Spatially ordered transcription of regulatory DNA in the bithorax complex of *Drosophila*. *Development* 107: 321–329.
- Bae E, Calhoun VC, Levine M, Lewis EB, Drewell RA (2002) Characterization of the intergenic RNA profile at *abdominal-A* and *-B* in the *Drosophila* bithorax complex. *Proc Natl Acad Sci U S A* 99: 16847–16852.
- Schmitt CH, Prestel M, Paro R (2005) Intergenic transcription through a polycomb group response element counteracts silencing. *Genes Dev* 19: 697–708.
- Petruk S, Sedkov Y, Riley KM, Hodgson J, Schweisguth F, et al. (2006) Transcription of bxd noncoding RNAs promoted by trithorax represses Ubx in cis by transcriptional interference. *Cell* 127: 1209–1221.
- Bender W (2008) MicroRNAs in the *Drosophila* bithorax complex. *Genes Dev* 22: 14–19.
- Stark A, Bushati N, Jan CH, Kheradpour P, Hodges E, et al. (2008) A single Hox locus in *Drosophila* produces functional microRNAs from opposite DNA strands. *Genes Dev* 22: 8–13.
- Tyler DM, Okamura K, Chung WJ, Hagen JW, Berezikov E, et al. (2008) Functionally distinct regulatory RNAs generated by bidirectional transcription and processing of microRNA loci. *Genes Dev* 22: 26–36.
- Bender W, Fitzgerald DP (2002) Transcription activates repressed domains in the *Drosophila* bithorax complex. *Development* 129: 4923–4930.
- Rank G, Prestel M, Paro R (2002) Transcription through intergenic chromosomal memory elements of the *Drosophila* bithorax complex correlates with an epigenetic switch. *Mol Cell Biol* 22: 8026–8034.
- Zhou J, Ashe H, Burks C, Levine M (1999) Characterization of the transvection mediating region of the *Abdominal-B* locus in *Drosophila*. *Development* 126: 3057–3065.
- Karch F, Bender W, Weiffenbach B (1990) abdA expression in *Drosophila* embryos. *Genes Dev* 4: 1573–1587.
- Harding K, Wedeen C, McGinnis W, Levine M (1985) Spatially regulated expression of homeotic genes in *Drosophila*. *Science* 229: 1236–1242.
- Duboule D (1991) Patterning in the vertebrate limb. *Curr Opin Genet Dev* 1: 211–216.
- Zavortink M, Sakonju S (1989) The morphogenetic and regulatory functions of the *Drosophila* *Abdominal-B* gene are encoded in overlapping RNAs transcribed from separate promoters. *Genes Dev* 3: 1969–1981.
- Enderle D, Beisel C, Stadler MB, Gerstung M, Athri P, et al. Polycomb preferentially targets stalled promoters of coding and noncoding transcripts. *Genome Res* 21: 216–226.
- Barges S, Mihaly J, Galloni M, Hagstrom K, Müller M, et al. (2000) The *Fab-8* boundary defines the distal limit of the bithorax complex *iab-7* domain and insulates *iab-7* from initiation elements and a PRE in the adjacent *iab-8* domain. *Development* 127: 779–790.
- Kondo T, Plaza S, Zanet J, Benrabah E, Valenti P, et al. (2010) Small peptides switch the transcriptional activity of Shavenbaby during *Drosophila* embryogenesis. *Science* 329: 336–339.

Author Contributions

Conceived and designed the experiments: MG RKM HG SS KAE FK WB. Performed the experiments: MG HG JJCA SS KAE WB. Analyzed the data: MG RKM HG SS KAE FK WB. Contributed reagents/materials/analysis tools: MG JJCA RKM HG SS KAE FK WB. Wrote the paper: MG RKM KAE FK WB.

- Galindo MI, Pueyo JI, Fouix S, Bishop SA, Couso JP (2007) Peptides encoded by short ORFs control development and define a new eukaryotic gene family. *PLoS Biol* 5: e106. doi:10.1371/journal.pbio.0050106.
- Lin H, Spradling AC (1993) Germline stem cell division and egg chamber development in transplanted *Drosophila* germaria. *Dev Biol* 159: 140–152.
- Brand AH, Perrimon N (1993) Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118: 401–415.
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, et al. (2007) Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 129: 1311–1323.
- Hopmann R, Duncan D, Duncan I (1995) Transvection in the *iab-5,6,7* region of the bithorax complex of *Drosophila*: homology independent interactions in trans. *Genetics* 139: 815–833. Issn: 0016–6731.
- Shearwin KE, Callen BP, Egan JB (2005) Transcriptional interference—a crash course. *Trends Genet* 21: 339–345.
- Penny GD, Kay GF, Sheardown SA, Rastan S, Brockdorff N (1996) Requirement for Xist in X chromosome inactivation. *Nature* 379: 131–137.
- Jeon Y, Lee JT (2011) YY1 tethers Xist RNA to the inactive X nucleation center. *Cell* 146: 119–133.
- Halic M, Moazed D (2010) Dicer-independent primal RNAs trigger RNAi and heterochromatin formation. *Cell* 140: 504–516.
- Wierzbicki AT, Haag JR, Pikaard CS (2008) Noncoding transcription by RNA polymerase Pol IVb/Pol V mediates transcriptional silencing of overlapping and adjacent genes. *Cell* 135: 635–648.
- Hagstrom K, Muller M, Schedl P (1997) A *Polycomb* and GAGA dependent silencer adjoins the *Fab-7* boundary in the *Drosophila* bithorax complex. *Genetics* 146: 1365–1380.
- Mihaly J, Hogga I, Gausz J, Gyurkovics H, Karch F (1997) In situ dissection of the *Fab-7* region of the bithorax complex into a chromatin domain boundary and a Polycomb-response element. *Development* 124: 1809–1820.
- Greger IH, Proudfoot NJ (1998) Poly(A) signals control both transcriptional termination and initiation between the tandem GAL10 and GAL7 genes of *Saccharomyces cerevisiae*. *EMBO J* 17: 4771–4779.
- Martens JA, Wu PY, Winston F (2005) Regulation of an intergenic transcript controls adjacent gene transcription in *Saccharomyces cerevisiae*. *Genes Dev* 19: 2695–2704.
- Buratowski S (2005) Connections between mRNA 3' end processing and transcription termination. *Curr Opin Cell Biol* 17: 257–261.
- Karch F, Weiffenbach B, Peifer M, Bender W, Duncan I, et al. (1985) The abdominal region of the bithorax complex. *Cell* 43: 81–96.
- Celniker SE, Sharma S, Keelan DJ, Lewis EB (1990) The molecular genetics of the bithorax complex of *Drosophila*: cis-regulation in the *Abdominal-B* domain. *Embo J* 9: 4277–4286.
- Gyurkovics H, Gausz J, Kummer J, Karch F (1990) A new homeotic mutation in the *Drosophila* bithorax complex removes a boundary separating two domains of regulation. *Embo J* 9: 2579–2585. Issn: 0261-4189.
- Mihaly J, Barges S, Sipos L, Maeda R, Cleard F, et al. (2006) Dissecting the regulatory landscape of the *Abd-B* gene of the bithorax complex. *Development* 133: 2983–2993.
- Nagaso H, Murata T, Day N, Yokoyama KK (2001) Simultaneous detection of RNA and protein by in situ hybridization and immunological staining. *J Histochem Cytochem* 49: 1177–1182.
- Karch F, Galloni M, Sipos L, Gausz J, Gyurkovics H, et al. (1994) Mcp and *Fab-7*: molecular analysis of putative boundaries of cis-regulatory domains in the bithorax complex of *Drosophila melanogaster*. *Nucleic Acids Res* 22: 3138–3146. Issn: 0305–1048.
- Holohan EE, Kwong C, Adryan B, Bartkuhn M, Herold M, et al. (2007) CTCF genomic binding sites in *Drosophila* and the organization of the bithorax complex. *PLoS Genet* 3: e112. doi:10.1371/journal.pgen.0030112.
- Cavener DR, Ray SC (1991) Eukaryotic start and stop translation sites. *Nucleic Acids Res* 19: 3185–3192.