

Illinois State University

ISU ReD: Research and eData

Theses and Dissertations

7-16-2021

On the Spectrum Problem for a Class of 3-Uniform Hypergraphs With 5 Edges

Kaitlin Elizabeth Shoukry

Illinois State University, Keshouk@ilstu.edu

Follow this and additional works at: <https://ir.library.illinoisstate.edu/etd>

Recommended Citation

Shoukry, Kaitlin Elizabeth, "On the Spectrum Problem for a Class of 3-Uniform Hypergraphs With 5 Edges" (2021). *Theses and Dissertations*. 1469.

<https://ir.library.illinoisstate.edu/etd/1469>

This Thesis is brought to you for free and open access by ISU ReD: Research and eData. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of ISU ReD: Research and eData. For more information, please contact ISUREd@ilstu.edu.

ON THE SPECTRUM PROBLEM FOR A CLASS OF
3-UNIFORM HYPERGRAPHS WITH 5 EDGES

KAITLIN ELIZABETH SHOUKRY

22 Pages

The complete 3-uniform hypergraph of order v has a set V of size v as its vertex set and the set of all 3-element subsets of V as its edge set. The degree of a vertex is the number of edges in its edge set that contain it. We consider a class of 3-uniform hypergraphs with 5 edges and 10 vertices such that: every vertex has degree either 1 or 2 and any two edges intersect in at most one vertex. There are 5 such hypergraphs. For $k \in \{1, 2, 3, 4, 5\}$, let H_k denote the hypergraphs with vertex set $\{v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}\}$ and edge sets $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_4, v_6\}, \{v_3, v_7, v_8\}, \{v_5, v_9, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_4, v_6\}, \{v_3, v_7, v_8\}, \{v_7, v_9, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_6, v_7\}, \{v_3, v_8, v_9\}, \{v_4, v_6, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_6, v_7\}, \{v_4, v_8, v_9\}, \{v_6, v_8, v_{10}\}\}$, and $\{\{v_1, v_2, v_3\}, \{v_3, v_4, v_5\}, \{v_5, v_6, v_7\}, \{v_7, v_8, v_9\}, \{v_9, v_{10}, v_0\}\}$, respectively. We give necessary and sufficient conditions for the existence of a decomposition of the complete 3-uniform hypergraph of order v into isomorphic copies of each H_k .

KEYWORDS: 3-Uniform Hypergraphs, Hypergraph Decompositions

ON THE SPECTRUM PROBLEM FOR A CLASS OF
3-UNIFORM HYPERGRAPHS WITH 5 EDGES

KAITLIN ELIZABETH SHOUKRY

A Thesis Submitted in Partial
Fulfillment of the Requirements
for the Degree of

MASTER OF SCIENCE

Department of Mathematics

ILLINOIS STATE UNIVERSITY

2021

© 2021 Kaitlin Elizabeth Shoukry

ON THE SPECTRUM PROBLEM FOR A CLASS OF
3-UNIFORM HYPERGRAPHS WITH 5 EDGES

KAITLIN ELIZABETH SHOUKRY

COMMITTEE MEMBERS:

Saad I. El-Zanati, Chair

Ryan C. Bunge

Papa A. Sissokho

ACKNOWLEDGMENTS

Part of this work was initiated in MAT461, *Advanced Topics in Discrete Mathematics*, taught by Professor Saad El-Zanati (assisted by Dr. Ryan Bunge) in Spring 2019. The template used for this work was shared by Drs. El-Zanati and Bunge.

K. E. S.

CONTENTS

	Page
ACKNOWLEDGMENTS	i
CONTENTS	ii
FIGURES	iii
CHAPTER I: INTRODUCTION	1
CHAPTER II: NOTATION AND TERMINOLOGY	6
CHAPTER III: EXAMPLES OF H_k -DECOMPOSITIONS	7
CHAPTER IV: MAIN RESULTS	19
REFERENCES	21

FIGURES

Figure		Page
1.	H_1 denoted $H_1[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$	4
2.	H_2 denoted $H_2[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$	4
3.	H_3 denoted $H_3[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$	4
4.	H_4 denoted $H_4[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$	4
5.	H_5 denoted $H_4[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$	5

CHAPTER I: INTRODUCTION

A *graph* G is an ordered pair $(V(G), E(G))$, where $V(G)$ is a finite set of elements called the *vertices* of G and $E(G)$ is a set of 2-element subsets of $V(G)$, called the *edges* of G . If $e = \{u, v\}$ is an edge in $E(G)$, then we say edge e is *incident* with vertices u and v and call u and v the *end-vertices* of e . In this case, we also say vertices u and v are *incident* with edge e . Two vertices u and v in $V(G)$ are *adjacent* in G if $\{u, v\} \in E(G)$. Similarly, edges e and e' are *adjacent* in G if e and e' share a common end-vertex. The *degree* of a vertex $v \in V(G)$ is the number of edges in $E(G)$ that contain v . We call $|V(G)|$ the *order* of G and $|E(G)|$ its *size*.

Two graphs $G = (V(G), E(G))$ and $G' = (V(G'), E(G'))$ are said to be *isomorphic* if there exists a one-to-one and onto map $f: V(G) \mapsto V(G')$ that preserves adjacency. Thus in this case, two vertices u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in G' .

A commonly studied problem in combinatorics concerns decompositions of graphs into edge-disjoint subgraphs. A *decomposition* of a graph K is a set $\Delta = \{G_1, G_2, \dots, G_s\}$ of pairwise edge-disjoint subgraphs of K such that $E(G_1) \cup E(G_2) \cup \dots \cup E(G_s) = E(K)$. If each element of Δ is isomorphic to a fixed graph G , then Δ is called a *G -decomposition* of K . A G -decomposition of K_v is also known as a *G -design of order v* . A K_k -design of order v is an *$S(2, k, v)$ -design* or a *Steiner system*. An $S(2, k, v)$ -design is also known as a *balanced incomplete block design of index 1* or a *$(v, k, 1)$ -BIBD*. The problem of determining all v for which there exists a G -design of order v is of special interest (see [1] for a survey).

The notion of decompositions of graphs naturally extends to decompositions of uniform hypergraphs. A *hypergraph* H consists of a finite nonempty set V of *vertices* and a set $E = \{e_1, e_2, \dots, e_m\}$ of nonempty subsets of V called *hyperedges*. If for each $e \in E$ we have $|e| = t$, then H is said to be *t -uniform*. Thus graphs are 2-uniform hypergraphs. The complete t -uniform hypergraph on the vertex set V has the set of all t -element subsets of V as its edge set and is denoted by $K_V^{(t)}$. If $v = |V|$, then $K_v^{(t)}$ is called the *complete t -uniform*

hypergraph of order v and is used to denote any hypergraph isomorphic to $K_V^{(t)}$.

A *decomposition* of a hypergraph K is a set $\Delta = \{H_1, H_2, \dots, H_s\}$ of pairwise edge-disjoint subgraphs of K such that $E(H_1) \cup E(H_2) \cup \dots \cup E(H_s) = E(K)$. If each element H_i of Δ is isomorphic to a fixed hypergraph H , then each H_i is called an *H-block*, and Δ is called an *H-decomposition* of K . If there exists an *H-decomposition* of K , then we may simply state that *H decomposes K*. An *H-decomposition* of the complete t -uniform hypergraph of order v is also called an *H-design of order v*. The problem of determining all v for which there exists an *H-design* of order v is called the *spectrum problem for H-designs*.

A $K_k^{(t)}$ -design of order v is a generalization of Steiner systems and is equivalent to an $S(t, k, v)$ -design. A summary of results on $S(t, k, v)$ -designs appears in [8]. Keevash [14] has recently shown that for all t and k the obvious necessary conditions for the existence of an $S(t, k, v)$ -design are sufficient for sufficiently large values of v . Similar results were obtained by Glock, Kühn, Lo, and Osthus [9, 10] and extended to include the corresponding asymptotic results for *H-designs* of order v for all uniform hypergraphs H . These results for t -uniform hypergraphs mirror the celebrated results of Wilson [19] for graphs. Although these asymptotic results assure the existence of *H-designs* for sufficiently large values of v for any uniform hypergraph H , the spectrum problem has been settled for very few hypergraphs of uniformity larger than 2.

In the study of graph decompositions, a fair amount of the focus has been on G -decompositions of K_v where G is a graph with a relatively small number of edges (see [1] and [6] for known results). Some authors have investigated the corresponding problem for 3-uniform hypergraphs. For example, in [4], the spectrum problem is settled for all 3-uniform hypergraphs on 4 or fewer vertices. More recently, the spectrum problem was settled in [5] for all 3-uniform hypergraphs with at most 6 vertices and at most 3 edges. In [5], they also settle the spectrum problem for the 3-uniform hypergraph of order 6 whose edges form the lines of the Pasch configuration. Authors have also considered *H-designs*

where H is a 3-uniform hypergraph whose edge set is defined by the faces of a regular polyhedron. Let T , O , and I denote the tetrahedron, the octahedron, and the icosahedron hypergraphs, respectively. The hypergraph T is the same as $K_4^{(3)}$, and its spectrum was settled in 1960 by Hanani [11]. In another paper [12], Hanani settled the spectrum problem for O -designs and gave necessary conditions for the existence of I -designs.

Perhaps the best known general result on decompositions of complete t -uniform hypergraphs is Baranyai's result [3] on the existence of 1-factorizations of $K_{mt}^{(t)}$ for all positive integers m . There are, however, several articles on decompositions of complete t -uniform hypergraphs (see [2] and [17]) and of t -uniform t -partite hypergraphs (see [15] and [18]) into variations on the concept of a Hamilton cycle. There are also several results on decompositions of 3-uniform hypergraphs into structures known as Berge cycles with a given number of edges (see for example [13] and [16]). We note however that the Berge cycles in these decompositions are not required to be isomorphic.

In this work, we considered the spectrum problem for the class of 3-uniform hypergraphs with 5 edges and 10 vertices, where the minimum vertex degree is 1, the maximum vertex degree is 2, and any two edges intersect in at most one vertex. There are 5 such hypergraphs as shown in Figures 1–2 below. For $k \in \{1, 2, 3, 4, 5\}$, let H_k denote the hypergraphs with vertex set $\{v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}\}$ and edge sets $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_4, v_6\}, \{v_3, v_7, v_8\}, \{v_5, v_9, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_4, v_6\}, \{v_3, v_7, v_8\}, \{v_7, v_9, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_6, v_7\}, \{v_3, v_8, v_9\}, \{v_4, v_6, v_{10}\}\}$, $\{\{v_1, v_2, v_3\}, \{v_1, v_4, v_5\}, \{v_2, v_6, v_7\}, \{v_4, v_8, v_9\}, \{v_6, v_8, v_{10}\}\}$, and $\{\{v_1, v_2, v_3\}, \{v_3, v_4, v_5\}, \{v_5, v_6, v_7\}, \{v_7, v_8, v_9\}, \{v_9, v_{10}, v_0\}\}$, respectively.

The graph H_5 is known as a *loose 5-cycle*. It is shown in [7] that there exists an H_5 -decomposition of $K_v^{(3)}$ if and only if $v \equiv 0, 1$ or $2 \pmod{5}$, and $v \geq 10$. We settle the spectrum problem for the remaining 4 hypergraphs.




Figure 1: H_1 denoted $H_1[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$




Figure 2: H_2 denoted $H_2[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$




Figure 3: H_3 denoted $H_3[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$




Figure 4: H_4 denoted $H_4[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$




Figure 5: H_5 denoted $H_4[v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, v_{10}]$

CHAPTER II: NOTATION AND TERMINOLOGY

If a and b are integers, we define $[a, b]$ to be $\{r \in \mathbb{Z} : a \leq r \leq b\}$. Let \mathbb{Z}_n denote the group of integers modulo n .

We will often describe our hypergraphs by giving their edge set only. Since the hypergraphs we consider will never contain isolated vertices, this is enough to uniquely define them. The complete k -uniform hypergraph with vertex set V has the set of all k -element subsets of V as its edge set and is denoted by $K_V^{(k)}$. If $v = |V|$, then $K_v^{(k)}$ is used to denote any hypergraph isomorphic to $K_V^{(k)}$. If H' is a subhypergraph of H , then $H \setminus H'$ denotes the hypergraph obtained from H by deleting the edges of H' .

We need to define some notation for certain types of multipartite hypergraphs. Let U_1, U_2, \dots, U_m be pairwise disjoint sets. The hypergraph with vertex set $V = U_1 \cup U_2 \cup \dots \cup U_m$ and edge set consisting of all k -element subsets of V having at most one vertex in each of U_1, U_2, \dots, U_m is denoted by $K_{U_1, U_2, \dots, U_m}^{(k)}$. If $|U_i| = u_i$ for $i \in [1, m]$, we may use $K_{u_1, u_2, \dots, u_m}^{(k)}$ to denote any hypergraph that is isomorphic to $K_{U_1, U_2, \dots, U_m}^{(k)}$, and if $u_1 = u_2 = \dots = u_m = u$, then the notation $K_{m \times u}^{(k)}$ may be used instead of $K_{u_1, u_2, \dots, u_m}^{(k)}$.

For pairwise disjoint sets U_1, U_2, \dots, U_m , $1 \leq m \leq k - 1$, the hypergraph with vertex set $V = U_1 \cup U_2 \cup \dots \cup U_m$ and edge set consisting of all k -element subsets of V having at least one element in each of U_1, U_2, \dots, U_m is denoted by $L_{U_1, U_2, \dots, U_m}^{(k)}$. If $|U_i| = u_i$ for $i \in [1, m]$, we may use $L_{u_1, u_2, \dots, u_m}^{(k)}$ to denote any hypergraph that is isomorphic to $L_{U_1, U_2, \dots, U_m}^{(k)}$. If k_1, k_2, \dots, k_m are positive integers with $k_1 + k_2 + \dots + k_m = k$, then $L_{U_1, U_2, \dots, U_m}^{(k_1, k_2, \dots, k_m)}$ is the subgraph of $L_{U_1, U_2, \dots, U_m}^{(k)}$ where each hyperedge contains exactly k_i elements from each U_i . We define $L_{u_1, u_2, \dots, u_m}^{(k_1, k_2, \dots, k_m)}$ similarly.

CHAPTER III: EXAMPLES OF H_k -DECOMPOSITIONS

We give several examples of H_k -decompositions, $k \in \{1, 2, 3, 4\}$, that are used in proving our main result.

Example 1. Let $V\left(K_{10}^{(3)}\right) = \mathbb{Z}_8 \cup \{\infty_1, \infty_2\}$ and let

$$B_1 = \{H_1[2, 1, 4, \infty_2, 5, 3, 0, 6, 7, \infty_1], H_1[0, 5, 2, \infty_1, 3, \infty_2, 1, 6, 4, 7]\},$$

$$B'_1 = \{H_1[\infty_1, 0, 7, 4, 3, \infty_2, 5, 6, 1, 2], H_1[\infty_1, 1, 0, 5, 4, \infty_2, 6, 7, 2, 3],$$

$$H_1[\infty_1, 2, 1, 6, 5, \infty_2, 7, 0, 3, 4], H_1[\infty_1, 3, 2, 7, 6, \infty_2, 0, 1, 4, 5],$$

$$H_1[\infty_2, 0, 7, 4, 3, \infty_1, 1, 2, 5, 6], H_1[\infty_2, 1, 0, 5, 4, \infty_1, 2, 3, 6, 7],$$

$$H_1[\infty_2, 2, 1, 6, 5, \infty_1, 3, 4, 7, 0], H_1[\infty_2, 3, 2, 7, 6, \infty_1, 4, 5, 0, 1]\},$$

$$B_2 = \{H_2[5, 2, 0, 7, 6, 1, 4, 3, \infty_1, \infty_2], H_2[6, 0, 4, 3, 2, \infty_1, 1, 7, 5]\},$$

$$B'_2 = \{H_2[4, 1, 2, \infty_2, 5, 0, 7, 6, 3, \infty_1], H_2[5, 2, 3, \infty_2, 6, 1, 0, 7, 4, \infty_1],$$

$$H_2[6, 3, 4, \infty_2, 7, 2, 1, 0, 5, \infty_1], H_2[7, 4, 5, \infty_2, 0, 3, 2, 1, 6, \infty_1],$$

$$H_2[5, 0, 6, \infty_1, 4, 1, 3, 2, 7, \infty_2], H_2[6, 1, 7, \infty_1, 5, 2, 4, 3, 0, \infty_2],$$

$$H_2[7, 2, 0, \infty_1, 6, 3, 5, 4, 1, \infty_2], H_2[0, 3, 1, \infty_1, 7, 4, 6, 5, 2, \infty_2]\},$$

$$B_3 = \{H_3[0, \infty_1, 3, 2, \infty_2, 4, 1, 7, 5, 6], H_3[0, 5, 2, 1, 7, 6, \infty_1, \infty_2, 4, 3]\},$$

$$B'_3 = \{H_3[\infty_1, 1, 0, 4, 5, 3, 2, 6, \infty_2, 7], H_3[\infty_1, 2, 1, 5, 6, 4, 3, 7, \infty_2, 0],$$

$$H_3[\infty_1, 3, 2, 6, 7, 5, 4, 0, \infty_2, 1], H_3[\infty_1, 4, 3, 7, 0, 6, 5, 1, \infty_2, 2],$$

$$H_3[\infty_2, 0, 1, 5, 4, 6, 3, 7, \infty_1, 2], H_3[\infty_2, 1, 2, 6, 5, 7, 4, 0, \infty_1, 3],$$

$$H_3[\infty_2, 2, 3, 7, 6, 0, 5, 1, \infty_1, 4], H_3[\infty_2, 3, 4, 0, 7, 1, 6, 2, \infty_1, 5]\},$$

$$B_4 = \{H_4[\infty_2, 0, 2, 3, 6, \infty_1, 5, 4, 7, 1], H_4[6, 2, 0, 4, 5, 7, 1, \infty_1, \infty_2, 3]\},$$

$$B'_4 = \{H_4[1, 7, 6, 4, 5, \infty_1, 0, 2, \infty_2, 3], H_4[2, 0, 7, 5, 6, \infty_1, 1, 3, \infty_2, 4],$$

$$H_4[3, 1, 0, 6, 7, \infty_1, 2, 4, \infty_2, 5], H_4[4, 2, 1, 7, 0, \infty_1, 3, 5, \infty_2, 6],$$

$$H_4[5, 3, 2, 0, 1, \infty_2, 4, 6, \infty_1, 7], H_4[6, 4, 3, 1, 2, \infty_2, 5, 7, \infty_1, 0],$$

$$H_4[7, 5, 4, 2, 3, \infty_2, 6, 0, \infty_1, 1], H_4[0, 6, 5, 3, 4, \infty_2, 7, 1, \infty_1, 2]\}.$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{10}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $\infty_i \mapsto \infty_i$, for $i \in \{1, 2\}$ and $j \mapsto j + 1 \pmod{8}$ along with the H_k -blocks in B'_k .

Example 2. Let $V(K_{11}^{(3)}) = \mathbb{Z}_{11}$ and let

$$B_1 = \{H_1[8, 3, 0, 10, 1, 4, 6, 7, 2, 5], H_1[4, 0, 7, 10, 6, 2, 9, 1, 3, 8], H_1[2, 0, 7, 10, 3, 5, 8, 9, 1, 4]\},$$

$$B_2 = \{H_2[3, 8, 0, 5, 1, 6, 9, 10, 2, 7], H_2[6, 0, 5, 2, 7, 8, 3, 10, 1, 4], H_2[4, 7, 0, 1, 5, 8, 9, 3, 10, 2]\},$$

$$B_3 = \{H_3[5, 4, 10, 6, 9, 1, 0, 3, 8, 7], H_3[1, 10, 8, 2, 6, 0, 3, 4, 7, 5], H_3[7, 0, 2, 3, 1, 9, 4, 5, 6, 8]\},$$

$$B_4 = \{H_4[5, 0, 6, 3, 8, 4, 1, 7, 9, 10], H_4[8, 0, 3, 4, 1, 5, 9, 6, 7, 2], H_4[9, 2, 0, 8, 7, 10, 4, 1, 5, 3]\}.$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{11}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $j \mapsto j + 1 \pmod{11}$.

Example 3. Let $V(K_{12}^{(3)}) = \mathbb{Z}_{11} \cup \{\infty\}$ and let

$$B_1 = \{H_1[5, 0, 2, 10, 6, 7, 4, 8, 9, \infty], H_1[0, 3, 8, 4, 1, 9, 7, 10, 6, \infty],$$

$$H_1[4, 7, 0, 8, 2, 3, 9, 10, 6, \infty], H_1[0, 2, 9, 1, \infty, 10, 4, 7, 3, 6]\},$$

$$B_2 = \{H_2[4, 0, 2, 10, 5, 7, 6, 3, 9, \infty], H_2[8, 3, 0, 6, 5, 1, 4, 9, 2, \infty],$$

$$H_2[7, 4, 0, 8, 5, 3, 9, 6, 2, \infty], H_2[0, 7, 2, 5, 4, 6, \infty, 8, 9, 10]\},$$

$$B_3 = \{H_3[0, 5, 2, 6, 10, 9, 1, 3, 7, \infty], H_3[3, 8, 0, 7, 2, 4, 1, 6, 10, \infty],$$

$$H_3[0, 4, 7, 9, 2, 3, 5, 6, 8, \infty], H_3[0, 1, \infty, 2, 8, 10, 5, 6, 7, 4]\},$$

$$B_4 = \{H_4[4, 0, 2, 8, 5, 6, 1, 10, \infty, 3], H_4[6, 3, 0, 7, 2, 8, 5, 9, \infty, 1],$$

$$H_4[4, 8, 0, 10, 6, 9, 5, 7, \infty, 1], H_4[2, 3, \infty, 7, 0, 5, 4, 1, 6, 8]\}.$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{12}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $\infty \mapsto \infty$ and $j \mapsto j + 1 \pmod{11}$.

Example 4. Let $V(K_{15}^{(3)}) = \mathbb{Z}_{13} \cup \{\infty_1, \infty_2\}$ and let

$$\begin{aligned}
B_1 = & \{H_1[0, 1, \infty_2, 12, \infty_1, 3, 4, 6, 7, 9], H_1[0, 1, 3, 12, 10, 6, 4, 8, 5, 9], \\
& H_1[0, 1, 4, 12, 9, 7, 5, 10, 3, 8], H_1[0, 3, 7, 6, 10, 1, 4, 12, 5, 8], \\
& H_1[0, 3, \infty_2, 10, \infty_1, 1, 2, 6, 7, 11], H_1[0, 5, \infty_2, 8, \infty_1, 11, 1, 7, 6, 12], \\
& H_1[0, 4, 8, 2, 6, 3, 1, 7, \infty_1, \infty_2]\}, \\
B_2 = & \{H_2[0, 1, 3, 12, 10, 6, 4, \infty_2, 5, \infty_1], H_2[0, 1, 4, 12, 9, 3, 6, \infty_2, 8, \infty_1], \\
& H_2[0, 1, 5, 12, 8, 4, 2, \infty_2, 6, \infty_1], H_2[0, 1, 6, 12, 7, 5, 2, \infty_2, 10, \infty_1], \\
& H_2[0, 3, 7, 6, 10, 9, 12, \infty_2, 2, \infty_1], H_2[0, 4, 9, 5, 8, 3, \infty_2, 2, 6, \infty_1], \\
& H_2[0, 6, 7, 8, 2, 3, 5, 1, 11, \infty_1]\}, \\
B_3 = & \{H_3[0, 3, 1, 12, 10, \infty_1, 4, 6, 8, \infty_2], H_3[0, 4, 1, 12, 9, \infty_1, 2, 6, 10, \infty_2], \\
& H_3[0, 5, 1, 12, 8, \infty_1, 4, 7, 10, \infty_2], H_3[0, 1, 6, 12, 7, \infty_1, 2, 5, 10, \infty_2], \\
& H_3[0, 5, 2, 8, 11, \infty_1, 4, 6, \infty_2, 9], H_3[0, 7, 2, 6, 11, \infty_1, 1, 4, \infty_2, 8], \\
& H_3[0, 6, 7, 2, 1, 9, 11, \infty_1, \infty_2, 5]\}, \\
B_4 = & \{H_4[0, 3, 1, 10, 12, 6, \infty_2, 4, 7, \infty_1], H_4[0, 4, 1, 9, 12, 6, \infty_2, 3, 5, \infty_1], \\
& H_4[0, 5, 1, 8, 12, 6, \infty_2, 4, 9, \infty_1], H_4[0, 5, 2, 11, 8, 12, \infty_2, 1, 3, \infty_1], \\
& H_4[0, 6, 1, 7, 12, 2, \infty_2, 5, 9, \infty_1], H_4[0, 6, 2, 7, 11, 1, \infty_2, 3, 9, \infty_1], \\
& H_4[0, 2, 7, 10, 6, 5, 9, \infty_1, \infty_2, 12]\}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{15}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $\infty_i \mapsto \infty_i$, for $i \in \{1, 2\}$, and $j \mapsto j + 1 \pmod{13}$.

Example 5. Let $V(K_{16}^{(3)}) = \mathbb{Z}_{16}$ and let

$$\begin{aligned}
B_1 = & \{H_1[0, 1, 3, 15, 13, 8, 5, 9, 7, 11], H_1[0, 1, 4, 15, 12, 13, 6, 11, 5, 10], \\
& H_1[0, 5, 1, 11, 15, 8, 3, 6, 10, 13], H_1[0, 6, 1, 10, 15, 2, 4, 9, 7, 12], \\
& H_1[0, 7, 1, 9, 15, 8, 5, 10, 6, 11], H_1[0, 7, 3, 9, 13, 15, 4, 11, 5, 12], \\
& H_1[0, 3, 9, 13, 7, 5, 4, 15, 1, 11]\}, \\
B_2 = & \{H_2[0, 3, 1, 13, 15, 8, 9, 4, 2, 12], H_2[0, 4, 1, 12, 15, 8, 6, 3, 9, 13], \\
& H_2[0, 5, 1, 11, 15, 8, 9, 3, 4, 12], H_2[0, 6, 1, 10, 15, 8, 7, 3, 2, 9], \\
& H_2[0, 7, 1, 9, 15, 8, 3, 11, 5, 10], H_2[0, 4, 9, 12, 7, 5, 3, 13, 1, 10], \\
& H_2[0, 10, 3, 9, 13, 2, 15, 5, 1, 12]\}, \\
B_3 = & \{H_3[0, 3, 1, 13, 15, 8, 6, 9, 14, 4], H_3[0, 4, 1, 12, 15, 8, 6, 9, 13, 5], \\
& H_3[0, 5, 1, 11, 15, 8, 4, 7, 13, 6], H_3[0, 6, 1, 10, 15, 8, 2, 5, 12, 7], \\
& H_3[0, 7, 1, 9, 15, 8, 2, 5, 13, 12], H_3[0, 9, 4, 7, 12, 1, 3, 6, 15, 11], \\
& H_3[0, 4, 10, 2, 12, 13, 6, 14, 15, 3]\}, \\
B_4 = & \{H_4[0, 3, 1, 13, 15, 8, 5, 4, 9, 11], H_4[0, 1, 4, 15, 12, 8, 3, 7, 11, 13], \\
& H_4[0, 1, 5, 15, 11, 7, 3, 8, 14, 13], H_4[0, 1, 6, 15, 10, 9, 3, 2, 11, 13], \\
& H_4[0, 1, 7, 9, 15, 11, 4, 3, 13, 2], H_4[0, 1, 14, 13, 3, 12, 7, 1, 9, 4], \\
& H_4[0, 1, 15, 12, 4, 13, 8, 2, 11, 5]\}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{16}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map and $j \mapsto j + 1 \pmod{16}$.

Example 6. Let $V(K_{17}^{(3)}) = \mathbb{Z}_{17}$ and let

$$\begin{aligned}
B_1 = & \{H_1[0, 3, 1, 14, 16, 8, 4, 10, 7, 13], H_1[0, 4, 1, 13, 16, 9, 3, 7, 10, 14], \\
& H_1[0, 5, 1, 12, 16, 8, 3, 10, 9, 14], H_1[0, 6, 1, 11, 16, 9, 3, 12, 5, 14],
\end{aligned}$$

$$\begin{aligned}
& H_1[0, 7, 1, 10, 16, 5, 3, 11, 8, 13], H_1[0, 8, 1, 16, 9, 3, 5, 12, 2, 13], \\
& H_1[0, 16, 1, 9, 8, 13, 4, 11, 10, 15], H_1[0, 5, 12, 3, 14, 7, 8, 16, 2, 11]\}, \\
B_2 = & \{H_2[0, 3, 1, 14, 16, 6, 4, 13, 8, 15], H_2[0, 4, 1, 13, 16, 7, 6, 3, 2, 8], \\
& H_2[0, 5, 1, 12, 16, 8, 4, 14, 6, 15], H_2[0, 1, 6, 11, 16, 4, 8, 12, 5, 10], \\
& H_2[0, 1, 7, 16, 10, 3, 5, 12, 9, 15], H_2[0, 1, 8, 9, 16, 4, 6, 14, 2, 11], \\
& H_2[0, 1, 16, 9, 8, 5, 14, 6, 7, 12], H_2[0, 5, 12, 11, 6, 8, 16, 7, 1, 9]\}, \\
B_3 = & \{H_3[0, 3, 1, 14, 16, 6, 5, 9, 15, 4], H_3[0, 4, 1, 13, 16, 7, 6, 10, 15, 5], \\
& H_3[0, 5, 1, 12, 16, 8, 2, 7, 11, 6], H_3[0, 1, 6, 11, 16, 4, 5, 7, 14, 8], \\
& H_3[0, 1, 7, 16, 10, 3, 4, 8, 15, 2], H_3[0, 1, 8, 9, 16, 4, 3, 5, 15, 10], \\
& H_3[0, 1, 16, 9, 8, 5, 7, 10, 12, 14], H_3[0, 5, 12, 11, 6, 8, 9, 13, 16, 10]\}, \\
B_4 = & \{H_4[0, 3, 1, 14, 16, 8, 5, 4, 9, 12], H_4[0, 4, 1, 13, 16, 6, 10, 9, 15, 11], \\
& H_4[0, 1, 5, 16, 12, 9, 3, 8, 15, 13], H_4[0, 1, 6, 16, 11, 8, 3, 9, 13, 14], \\
& H_4[0, 1, 7, 16, 10, 11, 3, 4, 13, 14], H_4[0, 1, 8, 16, 9, 10, 5, 2, 14, 13], \\
& H_4[0, 4, 10, 13, 7, 5, 3, 1, 14, 15], H_4[0, 5, 12, 2, 15, 1, 13, 4, 9, 10]\}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{17}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map and $j \mapsto j + 1 \pmod{17}$.

Example 7. Let $V(L_{5,5}^{(3)}) = \mathbb{Z}_{10}$ with vertex partition $\{\{0, 2, 4, 6, 8\}, \{1, 3, 5, 7, 9\}\}$ and let

$$\begin{aligned}
B_1 &= \{H_1[5, 3, 2, 7, 8, 0, 4, 9, 1, 6], H_1[0, 4, 1, 6, 9, 5, 2, 7, 3, 8]\}, \\
B_2 &= \{H_2[0, 7, 3, 2, 5, 6, 8, 4, 1, 9], H_2[7, 2, 0, 6, 4, 3, 9, 1, 5, 8]\}, \\
B_3 &= \{H_3[0, 2, 7, 3, 8, 9, 4, 5, 6, 1], H_3[5, 2, 4, 6, 8, 1, 0, 3, 7, 9]\}, \\
B_4 &= \{H_4[0, 3, 7, 1, 9, 6, 4, 5, 8, 2], H_4[5, 0, 2, 9, 6, 7, 1, 3, 8, 4]\}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $L_{5,5}^{(3)}$ consists of the orbits of the

H_k -blocks in B_k under the action of the map $j \mapsto j + 1 \pmod{10}$.

Example 8. Let $V\left(L_{5,5}^{(3)} \cup K_{1,5,5}^{(3)}\right) = \mathbb{Z}_{10} \cup \{\infty\}$ with vertex partition $\{\{\infty\}, \{0, 2, 4, 6, 8\}, \{1, 3, 5, 7, 9\}\}$ and let

$$B_1 = \{H_1[0, 1, 3, 7, 9, 4, 2, 8, 6, \infty], H_1[2, 0, 5, 9, 4, 1, 6, \infty, 3, 8]\},$$

$$B'_2 = \{H_1[2, 8, 1, 3, 9, \infty, 0, 4, 5, 6], H_1[4, 0, 3, 5, 1, \infty, 2, 6, 7, 8], H_1[6, 2, 5, 7, 3, \infty, 4, 8, 9, 0], \\ H_1[8, 4, 7, 9, 5, \infty, 6, 0, 1, 2], H_1[0, 6, 9, 1, 7, \infty, 8, 2, 3, 4]\},$$

$$B_2 = \{H_2[0, 1, 3, 7, 9, 4, 8, 2, 5, \infty], H_2[2, 9, 4, 0, 5, 1, 8, 3, 7, \infty]\},$$

$$B'_2 = \{H_2[0, 1, 7, 6, 9, \infty, 4, 3, 5, 8], H_2[2, 3, 9, 8, 1, \infty, 6, 5, 7, 0], H_2[4, 5, 1, 0, 3, \infty, 8, 7, 9, 2], \\ H_2[6, 7, 3, 2, 5, \infty, 0, 9, 1, 4], H_2[8, 9, 5, 4, 7, \infty, 2, 1, 3, 6]\},$$

$$B_3 = \{H_3[2, 0, 5, 9, 4, 3, 1, 8, \infty, 7], H_3[0, 6, 1, 9, 4, 8, 2, 3, \infty, 7]\},$$

$$B'_3 = \{H_3[4, 0, 3, 5, 1, \infty, 6, 7, 8, 2], H_3[6, 2, 5, 7, 3, \infty, 8, 9, 0, 4], H_3[8, 4, 7, 9, 5, \infty, 0, 1, 2, 6], \\ H_3[0, 6, 9, 1, 7, \infty, 2, 3, 4, 8], H_3[2, 8, 1, 3, 9, \infty, 4, 5, 6, 0]\},$$

$$B_4 = \{H_4[0, 4, 1, 6, 9, 3, \infty, 2, 7, 8], H_4[0, 1, 3, 7, 9, 4, 8, 6, \infty, 2]\},$$

$$B'_4 = \{H_4[6, 1, 2, 5, 4, 0, 9, 3, 7, \infty], H_4[8, 3, 4, 7, 6, 2, 1, 5, 9, \infty], H_4[0, 5, 6, 9, 8, 4, 3, 7, 1, \infty], \\ H_4[2, 7, 8, 1, 0, 6, 5, 9, 3, \infty], H_4[4, 9, 0, 3, 2, 8, 7, 1, 5, \infty]\}.$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $L_{5,5}^{(3)} \cup K_{1,5,5}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $\infty \mapsto \infty$ and $j \mapsto j + 1 \pmod{10}$ along with the H_k -blocks in B'_k .

Example 9. Let $V\left(L_{5,5}^{(3)} \cup K_{2,5,5}^{(3)}\right) = \mathbb{Z}_{10} \cup \{\infty_1, \infty_2\}$ with vertex partition $\{\{\infty_1, \infty_2\},$

$\{0, 2, 4, 6, 8\}, \{1, 3, 5, 7, 9\}$ and let

$$\begin{aligned}
B_1 &= \{H_1[0, 1, 6, 4, 9, 7, 3, \infty_1, 8, \infty_2], H_1[5, 2, 0, 4, 7, 9, 3, \infty_2, 6, 8]\}, \\
B'_1 &= \{H_1[2, 8, 1, 3, 9, \infty_1, 0, 4, 5, 6], H_1[4, 0, 3, 5, 1, \infty_1, 2, 6, 7, 8], \\
&\quad H_1[6, 2, 5, 7, 3, \infty_1, 4, 8, 9, 0], H_1[8, 4, 7, 9, 5, \infty_1, 6, 0, 1, 2], \\
&\quad H_1[0, 6, 9, 1, 7, \infty_1, 8, 2, 3, 4], H_1[0, 1, \infty_1, 2, 3, 9, 5, 6, 8, \infty_2], \\
&\quad H_1[2, 3, \infty_1, 4, 5, 1, 7, 8, 0, \infty_2], H_1[4, 5, \infty_1, 6, 7, 3, 9, 0, 2, \infty_2], \\
&\quad H_1[6, 7, \infty_1, 8, 9, 5, 1, 2, 4, \infty_2], H_1[8, 9, \infty_1, 0, 1, 7, 3, 4, 6, \infty_2]\}, \\
B_2 &= \{H_2[0, 1, 3, 9, 7, 4, \infty_2, 2, 5, 8], H_2[3, 7, 0, 8, 1, 9, 5, 6, 2, \infty_1]\}, \\
B'_2 &= \{H_2[0, 1, 7, 6, 9, \infty_1, 4, 3, 5, 8], H_2[2, 3, 9, 8, 1, \infty_1, 6, 5, 7, 0], \\
&\quad H_2[4, 5, 1, 0, 3, \infty_1, 8, 7, 9, 2], H_2[6, 7, 3, 2, 5, \infty_1, 0, 9, 1, 4], \\
&\quad H_2[8, 9, 5, 4, 7, \infty_1, 2, 1, 3, 6], H_2[\infty_1, 0, 1, 5, 6, \infty_2, 2, 7, 3, 8], \\
&\quad H_2[\infty_1, 2, 3, 7, 8, \infty_2, 4, 9, 5, 0], H_2[\infty_1, 4, 5, 9, 0, \infty_2, 6, 1, 7, 2], \\
&\quad H_2[\infty_1, 6, 7, 1, 2, \infty_2, 8, 3, 9, 4], H_2[\infty_1, 8, 9, 3, 4, \infty_2, 0, 5, 1, 6]\}, \\
B_3 &= \{H_3[0, 1, 3, 7, 9, 2, 5, 8, \infty_1, 6], H_3[1, 0, \infty_2, 3, 6, 5, 2, 7, 8, 9]\}, \\
B'_3 &= \{H_3[4, 0, 3, 5, 1, \infty_1, 6, 7, 8, 2], H_3[6, 2, 5, 7, 3, \infty_1, 8, 9, 0, 4], \\
&\quad H_3[8, 4, 7, 9, 5, \infty_1, 0, 1, 2, 6], H_3[0, 6, 9, 1, 7, \infty_1, 2, 3, 4, 8], \\
&\quad H_3[2, 8, 1, 3, 9, \infty_1, 4, 5, 6, 0], H_3[\infty_1, 1, 0, 5, 6, 2, 4, 9, \infty_2, 7], \\
&\quad H_3[\infty_1, 3, 2, 7, 8, 4, 6, 1, \infty_2, 9], H_3[\infty_1, 5, 4, 9, 0, 6, 8, 3, \infty_2, 1], \\
&\quad H_3[\infty_1, 7, 6, 1, 2, 8, 0, 5, \infty_2, 3], H_3[\infty_1, 9, 8, 3, 4, 0, 2, 7, \infty_2, 5]\}, \\
B_4 &= \{H_4[2, 0, 5, 4, 9, \infty_1, 6, 8, 7], H_4[0, 1, 6, 7, 3, \infty_2, 2, 8, 9, 4]\}, \\
B'_4 &= \{H_4[6, 1, 2, 5, 4, 0, 9, 3, 7, \infty_1], H_4[8, 3, 4, 7, 6, 2, 1, 5, 9, \infty_1], \\
&\quad H_4[0, 5, 6, 9, 8, 4, 3, 7, 1, \infty_1], H_4[2, 7, 8, 1, 0, 6, 5, 9, 3, \infty_1], \\
&\quad H_4[4, 9, 0, 3, 2, 8, 7, 1, 5, \infty_1], H_4[5, 9, 8, 1, 2, 0, 6, 3, \infty_2, 4]\},
\end{aligned}$$

$$H_4[7, 1, 0, 3, 4, 2, 8, 5, \infty_2, 6], H_4[9, 3, 2, 5, 6, 4, 0, 7, \infty_2, 8],$$

$$H_4[1, 5, 4, 7, 8, 6, 2, 9, \infty_2, 0], H_4[3, 7, 6, 9, 0, 8, 4, 1, \infty_2, 2]\}.$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $L_{5,5}^{(3)} \cup K_{2,5,5}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map $\infty_i \mapsto \infty_i$, for $i \in \{1, 2\}$, and $j \mapsto j + 1 \pmod{10}$ along with the H_k -blocks in B'_k .

Example 10. Let $V\left(K_{3,5,5}^{(3)}\right) = \mathbb{Z}_{10} \cup \{\infty_1, \infty_2, \infty_3\}$ with vertex partition $\{\{\infty_1, \infty_2, \infty_3\}, \{0, 2, 4, 6, 8\}, \{1, 3, 5, 7, 9\}\}$ and let

$$B_1 = \{H_1[1, \infty_1, 0, 4, \infty_2, 5, 3, \infty_3, 2, 7], H_1[1, \infty_2, 0, 4, \infty_3, 5, 3, \infty_1, 2, 7],$$

$$H_1[1, \infty_3, 0, 4, \infty_1, 5, 3, \infty_2, 2, 7], H_1[9, \infty_1, 4, 2, \infty_2, 5, 3, \infty_3, 1, 6],$$

$$H_1[9, \infty_2, 4, 2, \infty_3, 5, 3, \infty_1, 1, 6], H_1[9, \infty_3, 4, 2, \infty_1, 5, 3, \infty_2, 1, 6],$$

$$H_1[6, \infty_1, 3, 7, \infty_2, 8, 2, \infty_3, 0, 5], H_1[6, \infty_2, 3, 7, \infty_3, 8, 2, \infty_1, 0, 5],$$

$$H_1[6, \infty_3, 3, 7, \infty_1, 8, 2, \infty_2, 0, 5], H_1[8, 9, \infty_1, \infty_2, 1, 0, 4, 7, 2, \infty_3],$$

$$H_1[8, 9, \infty_2, \infty_3, 1, 0, 4, 7, 2, \infty_1], H_1[8, 9, \infty_3, \infty_1, 1, 0, 4, 7, 2, \infty_2],$$

$$H_1[5, \infty_1, 8, 6, \infty_2, 9, 3, \infty_3, 0, 7], H_1[5, \infty_2, 8, 6, \infty_3, 9, 3, \infty_1, 0, 7],$$

$$H_1[5, \infty_3, 8, 6, \infty_1, 9, 3, \infty_2, 0, 7]\},$$

$$B_2 = \{H_2[\infty_2, 0, 5, 3, 4, \infty_1, \infty_3, 8, 2, 9], H_2[\infty_3, 0, 5, 3, 4, \infty_2, \infty_1, 8, 2, 9],$$

$$H_2[\infty_1, 0, 5, 3, 4, \infty_3, \infty_2, 8, 2, 9], H_2[\infty_2, 1, 6, 4, 5, \infty_1, \infty_3, 9, 7, 8],$$

$$H_2[\infty_1, 1, 6, 4, 5, \infty_2, \infty_1, 9, 7, 8], H_2[\infty_3, 1, 6, 4, 5, \infty_3, \infty_2, 9, 7, 8],$$

$$H_2[\infty_2, 2, 7, 5, 6, \infty_1, \infty_3, 0, 8, 9], H_2[\infty_3, 2, 7, 5, 6, \infty_2, \infty_1, 0, 8, 9],$$

$$H_2[\infty_1, 2, 7, 5, 6, \infty_3, \infty_2, 0, 8, 9], H_2[1, \infty_2, 8, 0, \infty_1, 9, \infty_3, 3, 6, 7],$$

$$H_2[1, \infty_3, 8, 0, \infty_2, 9, \infty_1, 3, 6, 7], H_2[1, \infty_1, 8, 0, \infty_3, 9, \infty_2, 3, 6, 7],$$

$$H_2[2, 3, \infty_1, \infty_2, 1, 6, 4, 9, 7, \infty_3], H_2[2, 3, \infty_2, \infty_3, 1, 6, 4, 9, 7, \infty_1],$$

$$H_2[2, 3, \infty_3, \infty_1, 1, 6, 4, 9, 7, \infty_2]\},$$

$$\begin{aligned}
B_3 = \{ & H_3[1, \infty_1, 0, 4, \infty_2, 7, 8, 9, \infty_3, 5], H_3[1, \infty_2, 0, 4, \infty_3, 7, 8, 9, \infty_1, 5], \\
& H_3[1, \infty_3, 0, 4, \infty_1, 7, 8, 9, \infty_2, 5], H_3[2, \infty_1, 9, 5, \infty_2, 4, 8, 1, \infty_3, 6], \\
& H_3[2, \infty_2, 9, 5, \infty_3, 4, 8, 1, \infty_1, 6], H_3[2, \infty_3, 9, 5, \infty_1, 4, 8, 1, \infty_2, 6], \\
& H_3[0, \infty_1, 3, 7, \infty_2, 2, 5, 8, \infty_3, 6], H_3[0, \infty_2, 3, 7, \infty_3, 2, 5, 8, \infty_1, 6], \\
& H_3[0, \infty_3, 3, 7, \infty_1, 2, 5, 8, \infty_2, 6], H_3[2, \infty_1, 1, 3, \infty_2, 4, 0, 9, \infty_3, 6], \\
& H_3[2, \infty_2, 1, 3, \infty_3, 4, 0, 9, \infty_1, 6], H_3[2, \infty_3, 1, 3, \infty_1, 4, 0, 9, \infty_2, 6], \\
& H_3[7, \infty_1, 6, 8, \infty_2, 3, 4, 9, \infty_3, 5], H_3[7, \infty_2, 6, 8, \infty_3, 3, 4, 9, \infty_1, 5], \\
& H_3[7, \infty_3, 6, 8, \infty_1, 3, 4, 9, \infty_2, 5] \}, \\
B_4 = \{ & H_4[0, 1, \infty_1, 7, \infty_3, \infty_2, 2, 3, 8, 4], H_4[0, 1, \infty_2, 7, \infty_1, \infty_3, 2, 3, 8, 4], \\
& H_4[0, 1, \infty_3, 7, \infty_2, \infty_1, 2, 3, 8, 4], H_4[2, 3, \infty_1, 9, \infty_3, \infty_2, 4, 5, 0, 6], \\
& H_4[2, 3, \infty_2, 9, \infty_1, \infty_3, 4, 5, 0, 6], H_4[2, 3, \infty_3, 9, \infty_2, \infty_1, 4, 5, 0, 6], \\
& H_4[4, 5, \infty_1, 1, \infty_3, \infty_2, 6, 7, 2, 8], H_4[4, 5, \infty_2, 1, \infty_1, \infty_3, 6, 7, 2, 8], \\
& H_4[4, 5, \infty_3, 1, \infty_2, \infty_1, 6, 7, 2, 8], H_4[6, 7, \infty_1, 3, \infty_3, \infty_2, 8, 9, 4, 0], \\
& H_4[6, 7, \infty_2, 3, \infty_1, \infty_3, 8, 9, 4, 0], H_4[6, 7, \infty_3, 3, \infty_2, \infty_1, 8, 9, 4, 0], \\
& H_4[8, 9, \infty_1, 5, \infty_3, \infty_2, 0, 1, 6, 2], H_4[8, 9, \infty_2, 5, \infty_1, \infty_3, 0, 1, 6, 2], \\
& H_4[8, 9, \infty_3, 5, \infty_2, \infty_1, 0, 1, 6, 2] \}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, B_k is an H_k -decomposition of $K_{3,5,5}^{(3)}$.

Example 11. Let $V(K_{4,5,5}^{(3)}) = \mathbb{Z}_{10} \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}$ with vertex partition $\{\{\infty_1, \infty_2, \infty_3, \infty_4\}, \{0, 2, 4, 6, 8\}, \{1, 3, 5, 7, 9\}\}$ and let

$$\begin{aligned}
B_1 = \{ & H_1[0, \infty_1, 1, 3, \infty_2, 6, 2, \infty_3, 4, 9], H_1[0, \infty_2, 1, 3, \infty_3, 6, 2, \infty_4, 4, 9], \\
& H_1[0, \infty_3, 1, 3, \infty_4, 6, 2, \infty_1, 4, 9], H_1[0, \infty_4, 1, 3, \infty_1, 6, 2, \infty_2, 4, 9], \\
& H_1[4, \infty_1, 1, 7, \infty_2, 0, 6, \infty_3, 2, 3], H_1[4, \infty_2, 1, 7, \infty_3, 0, 6, \infty_4, 2, 3], \\
& H_1[4, \infty_3, 1, 7, \infty_4, 0, 6, \infty_1, 2, 3], H_1[4, \infty_4, 1, 7, \infty_1, 0, 6, \infty_2, 2, 3],
\end{aligned}$$

$$\begin{aligned}
& H_1[9, 0, \infty_1, \infty_2, 8, 5, 3, 4, 7, \infty_3], H_1[9, 0, \infty_2, \infty_3, 8, 5, 3, 4, 7, \infty_4], \\
& H_1[9, 0, \infty_3, \infty_4, 8, 5, 3, 4, 7, \infty_1], H_1[9, 0, \infty_4, \infty_1, 8, 5, 3, 4, 7, \infty_2], \\
& H_1[5, 4, \infty_1, \infty_2, 8, 2, 0, 7, 3, \infty_3], H_1[5, 4, \infty_2, \infty_3, 8, 2, 0, 7, 3, \infty_4], \\
& H_1[5, 4, \infty_3, \infty_4, 8, 2, 0, 7, 3, \infty_1], H_1[5, 4, \infty_4, \infty_1, 8, 2, 0, 7, 3, \infty_2], \\
& H_1[7, 8, \infty_1, \infty_2, 2, 1, 5, 6, 9, \infty_3], H_1[7, 8, \infty_2, \infty_3, 2, 1, 5, 6, 9, \infty_4], \\
& H_1[7, 8, \infty_3, \infty_4, 2, 1, 5, 6, 9, \infty_1], H_1[7, 8, \infty_4, \infty_1, 2, 1, 5, 6, 9, \infty_2] \}, \\
B_2 = & \{ H_2[1, \infty_1, 0, 4, \infty_2, 5, \infty_3, 7, 8, 9], H_2[1, \infty_2, 0, 4, \infty_3, 5, \infty_4, 7, 8, 9], \\
& H_2[1, \infty_3, 0, 4, \infty_4, 5, \infty_1, 7, 8, 9], H_2[1, \infty_4, 0, 4, \infty_1, 5, \infty_2, 7, 8, 9], \\
& H_2[0, \infty_1, 5, 3, \infty_2, 8, 2, \infty_3, 7, \infty_4], H_2[0, \infty_2, 5, 3, \infty_3, 8, 2, \infty_4, 7, \infty_1], \\
& H_2[0, \infty_3, 5, 3, \infty_4, 8, 2, \infty_1, 7, \infty_2], H_2[0, \infty_4, 5, 3, \infty_1, 8, 2, \infty_2, 7, \infty_3], \\
& H_2[9, \infty_1, 2, 6, \infty_2, 7, \infty_3, 3, 1, 8], H_2[9, \infty_2, 2, 6, \infty_3, 7, \infty_4, 3, 1, 8], \\
& H_2[9, \infty_3, 2, 6, \infty_4, 7, \infty_1, 3, 1, 8], H_2[9, \infty_4, 2, 6, \infty_1, 7, \infty_2, 3, 1, 8], \\
& H_2[6, \infty_1, 1, 5, \infty_2, 8, \infty_3, 2, 4, 9], H_2[6, \infty_2, 1, 5, \infty_3, 8, \infty_4, 2, 4, 9], \\
& H_2[6, \infty_3, 1, 5, \infty_4, 8, \infty_1, 2, 4, 9], H_2[6, \infty_4, 1, 5, \infty_1, 8, \infty_2, 2, 4, 9], \\
& H_2[4, \infty_1, 3, 7, \infty_2, 8, \infty_3, 6, 0, 9], H_2[4, \infty_2, 3, 7, \infty_3, 8, \infty_4, 6, 0, 9], \\
& H_2[4, \infty_3, 3, 7, \infty_4, 8, \infty_1, 6, 0, 9], H_2[4, \infty_4, 3, 7, \infty_1, 8, \infty_2, 6, 0, 9] \}, \\
B_3 = & \{ H_3[1, \infty_1, 0, 4, \infty_2, 7, 8, 9, \infty_3, 5], H_3[1, \infty_2, 0, 4, \infty_3, 7, 8, 9, \infty_4, 5], \\
& H_3[1, \infty_3, 0, 4, \infty_4, 7, 8, 9, \infty_1, 5], H_3[1, \infty_4, 0, 4, \infty_1, 7, 8, 9, \infty_2, 5], \\
& H_3[2, \infty_1, 9, 5, \infty_2, 4, 8, 1, \infty_3, 6], H_3[2, \infty_2, 9, 5, \infty_3, 4, 8, 1, \infty_4, 6], \\
& H_3[2, \infty_3, 9, 5, \infty_4, 4, 8, 1, \infty_1, 6], H_3[2, \infty_4, 9, 5, \infty_1, 4, 8, 1, \infty_2, 6], \\
& H_3[0, \infty_1, 3, 7, \infty_2, 2, 5, 8, \infty_3, 6], H_3[0, \infty_2, 3, 7, \infty_3, 2, 5, 8, \infty_4, 6], \\
& H_3[0, \infty_3, 3, 7, \infty_4, 2, 5, 8, \infty_1, 6], H_3[0, \infty_4, 3, 7, \infty_1, 2, 5, 8, \infty_2, 6], \\
& H_3[2, \infty_1, 1, 3, \infty_2, 4, 0, 9, \infty_3, 6], H_3[2, \infty_2, 1, 3, \infty_3, 4, 0, 9, \infty_4, 6], \\
& H_3[2, \infty_3, 1, 3, \infty_4, 4, 0, 9, \infty_1, 6], H_3[2, \infty_4, 1, 3, \infty_1, 4, 0, 9, \infty_2, 6],
\end{aligned}$$

$$\begin{aligned}
& H_3[7, \infty_1, 6, 8, \infty_2, 3, 4, 9, \infty_3, 5], H_3[7, \infty_2, 6, 8, \infty_3, 3, 4, 9, \infty_4, 5], \\
& H_3[7, \infty_3, 6, 8, \infty_4, 3, 4, 9, \infty_1, 5], H_3[7, \infty_4, 6, 8, \infty_1, 3, 4, 9, \infty_2, 5] \}, \\
B_4 = & \{ H_4[1, \infty_1, 0, 6, \infty_2, 5, 2, 7, \infty_4, \infty_3], H_4[1, \infty_2, 0, 6, \infty_3, 5, 2, 7, \infty_1, \infty_4], \\
& H_4[1, \infty_3, 0, 6, \infty_4, 5, 2, 7, \infty_2, \infty_1], H_4[1, \infty_4, 0, 6, \infty_1, 5, 2, 7, \infty_3, \infty_2], \\
& H_4[2, \infty_1, 1, 7, \infty_2, 6, 3, 8, \infty_4, \infty_3], H_4[2, \infty_2, 1, 7, \infty_3, 6, 3, 8, \infty_1, \infty_4], \\
& H_4[2, \infty_3, 1, 7, \infty_4, 6, 3, 8, \infty_2, \infty_1], H_4[2, \infty_4, 1, 7, \infty_1, 6, 3, 8, \infty_3, \infty_2], \\
& H_4[3, \infty_1, 2, 8, \infty_2, 7, 4, 9, \infty_4, \infty_3], H_4[3, \infty_2, 2, 8, \infty_3, 7, 4, 9, \infty_1, \infty_4], \\
& H_4[3, \infty_3, 2, 8, \infty_4, 7, 4, 9, \infty_2, \infty_1], H_4[3, \infty_4, 2, 8, \infty_1, 7, 4, 9, \infty_3, \infty_2], \\
& H_4[4, \infty_1, 3, 9, \infty_2, 8, 5, 0, \infty_4, \infty_3], H_4[4, \infty_2, 3, 9, \infty_3, 8, 5, 0, \infty_1, \infty_4], \\
& H_4[4, \infty_3, 3, 9, \infty_4, 8, 5, 0, \infty_2, \infty_1], H_4[4, \infty_4, 3, 9, \infty_1, 8, 5, 0, \infty_3, \infty_2], \\
& H_4[5, \infty_1, 4, 0, \infty_2, 9, 6, 1, \infty_4, \infty_3], H_4[5, \infty_2, 4, 0, \infty_3, 9, 6, 1, \infty_1, \infty_4], \\
& H_4[5, \infty_3, 4, 0, \infty_4, 9, 6, 1, \infty_2, \infty_1], H_4[5, \infty_4, 4, 0, \infty_1, 9, 6, 1, \infty_3, \infty_2] \}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, B_k is an H_k -decomposition of $K_{4,5,5}^{(3)}$.

Example 12. Let $V(K_{5,5,5}^{(3)}) = \mathbb{Z}_{15}$ with vertex partition $\{\{0, 3, 6, 9, 12\}, \{1, 4, 7, 10, 13\}, \{2, 5, 8, 11, 14\}\}$ and let

$$\begin{aligned}
B_1 = & \{ H_1[2, 1, 0, 12, 10, 5, 7, 8, 9, 14], H_1[3, 2, 1, 13, 11, 6, 8, 9, 10, 0], \\
& H_1[4, 3, 2, 14, 12, 7, 9, 10, 11, 1], H_1[0, 11, 1, 10, 2, 6, 9, 14, 3, 13], \\
& H_1[13, 3, 8, 11, 9, 1, 4, 6, 5, 7] \}, \\
B_2 = & \{ H_2[2, 0, 1, 4, 6, 8, 11, 3, 7, 12, H_2[3, 1, 2, 5, 7, 9, 12, 4, 8, 13], \\
& H_2[4, 2, 3, 6, 8, 10, 13, 5, 9, 14], H_2[1, 8, 3, 9, 2, 13, 7, 14, 5, 12], \\
& H_2[1, 2, 6, 0, 5, 7, 4, 11, 9, 14] \},
\end{aligned}$$

$$\begin{aligned}
B_3 = & \{H_3[0, 1, 5, 14, 10, 9, 2, 7, 12, 3], H_3[1, 2, 6, 0, 11, 10, 5, 7, 9, 4], \\
& H_3[2, 3, 7, 1, 12, 11, 0, 4, 8, 5], H_3[0, 8, 1, 4, 2, 12, 5, 6, 13, 9], \\
& H_3[1, 2, 3, 0, 14, 13, 9, 10, 11, 7]\}, \\
B_4 = & \{H_4[0, 1, 5, 14, 10, 9, 2, 4, 6, 7], H_4[1, 2, 6, 0, 11, 10, 3, 5, 7, 8], \\
& H_4[2, 3, 7, 1, 12, 11, 4, 6, 8, 9], H_4[4, 0, 2, 12, 8, 10, 5, 9, 13, 14], \\
& H_4[4, 11, 0, 6, 2, 1, 3, 5, 7, 8]\}.
\end{aligned}$$

Then, for $k \in \{1, 2, 3, 4\}$, an H_k -decomposition of $K_{5,5,5}^{(3)}$ consists of the orbits of the H_k -blocks in B_k under the action of the map and $j \mapsto j + 3 \pmod{15}$.

CHAPTER IV: MAIN RESULTS

We begin by giving necessary conditions for the existence of an H_k -decomposition of $K_v^{(3)}$. An obvious necessary condition is that 5 must divide the number of edges in $K_v^{(3)}$, and thus we must have $v \equiv 0, 1, \text{ or } 2 \pmod{5}$. Also since H_k has 10 vertices, we must also have $n \geq 10$ for a non-trivial H_k -decomposition of $K_v^{(3)}$. Thus we have the following.

Lemma 1. *There exists an H_k -decomposition of $K_v^{(3)}$ only if $v \equiv 0, 1, \text{ or } 2 \pmod{5}$ and $v \geq 10$.*

We show that the above conditions are sufficient by showing how to construct H_k -decompositions of $K_v^{(3)}$ for all $v \equiv 0, 1, \text{ or } 2 \pmod{5}$ with $v \geq 10$. Our constructions are dependent on the many small examples given in Chapter III.

We begin by proving a lemma that is fundamental to our constructions.

Lemma 2. *For $r \in \{0, 1, 2\}$ and all positive integers x and y , there exists a decomposition of $K_{r,5x,5y}^{(3)} \cup L_{5x,5y}^{(3)}$ into copies of $K_{5,5,5}^{(3)}$ and $K_{r,5,5}^{(3)} \cup L_{5,5}^{(3)}$.*

Proof. Let $r \in \{0, 1, 2\}$ and let x and y be positive integers. The vertices of $K_{r,5x,5y}^{(3)} \cup L_{5x,5y}^{(3)}$ can be partitioned into sets V_i , W_j , and R where $1 \leq i \leq x$, $1 \leq j \leq y$, $|V_i| = 5 = |W_j|$, and $|R| = r$ such that every edge $\{a, b, c\}$ is of exactly one of the following types:

- Type 1: there exist i, j with $a \in R$, $b \in V_i$, and $c \in W_j$;
- Type 2: there exist i, j, k with $i \neq j$, $a \in V_i$, $b \in V_j$, and $c \in W_k$;
- Type 3: there exist i, j, k with $j \neq k$, $a \in V_i$, $b \in W_j$, and $c \in W_k$;
- Type 4: there exist i, j with $a, b \in V_i$ and $c \in W_j$; or
- Type 5: there exist i, j with $a \in V_i$ and $b, c \in W_j$.

For every choice of i and j we can put together the edges of Types 1, 4, and 5 to form a copy of $K_{r,5,5}^{(3)} \cup L_{5,5}^{(3)}$. For every choice of i, j , and k the edges of Types 2 and 3 form copies of $K_{5,5,5}^{(3)}$. Since all edges are accounted for by exactly one of the aforementioned choices of subscripts, we have the desired decomposition into copies of $K_{5,5,5}^{(3)}$ and $K_{r,5,5}^{(3)} \cup L_{5,5}^{(3)}$. \square

Theorem 3. *There exists an H_k -decomposition of $K_v^{(3)}$ if and only if $v \equiv 0, 1, \text{ or } 2 \pmod{5}$ and $v \geq 10$.*

Proof. The necessary conditions for the existence of an H_k -decomposition of $K_v^{(3)}$ are established in Lemma 1. Thus we need only to establish their sufficiency. Let $v = 5x + r$ where $x \geq 2$ and $r \in \{0, 1, 2\}$. We will consider two cases depending on the parity of x .

When x is even we can write $K_v^{(3)}$ as the edge-disjoint union of copies of $K_{10+r}^{(3)}$, $K_{r,10,10}^{(3)} \cup L_{10,10}^{(3)}$, and $K_{10,10,10}^{(3)}$, where the group of r vertices is common to every applicable copy. By Examples 1, 2, and 3 we have that an H_k -decomposition of $K_{10+r}^{(3)}$ exists. By Lemma 2 we have that $K_{r,10,10}^{(3)} \cup L_{10,10}^{(3)}$ can be decomposed into copies of $K_{5,5,5}^{(3)}$ and $K_{r,5,5}^{(3)} \cup L_{5,5}^{(3)}$. By Example 12 an H_k -decomposition of $K_{5,5,5}^{(3)}$ exists. When $r = 0$ $K_{0,5,5}^{(3)} \cup L_{5,5}^{(3)}$ is isomorphic to $L_{5,5}^{(3)}$, which admits an H_k -decomposition by Example 7. When $r \in \{1, 2\}$ an H_k -decomposition of $K_{r,5,5}^{(3)} \cup L_{5,5}^{(3)}$ exists by Examples 8 and 9. Finally, it is straightforward to see that $K_{10,10,10}^{(3)}$ can be decomposed into copies of $K_{5,5,5}^{(3)}$; thus, by Example 12 an H_k -decomposition of $K_{10,10,10}^{(3)}$ exists.

When x is odd the construction is similar, and we can write $K_v^{(3)}$ as the edge disjoint union of copies of $K_{15+r}^{(3)}$, $K_{10+r}^{(3)}$, $K_{r,15,10}^{(3)} \cup L_{15,10}^{(3)}$, $K_{r,10,10}^{(3)} \cup L_{10,10}^{(3)}$, $K_{15,10,10}^{(3)}$, and $K_{10,10,10}^{(3)}$, where again the group of r vertices is common to every applicable copy. There exist H_k -decompositions of each of these hypergraphs exist by using the ingredients listed in the previous case along with Examples 4, 5, and 6. □

REFERENCES

- [1] P. Adams, D. Bryant, and M. Buchanan, A survey on the existence of G -designs, *J. Combin. Des.* **16** (2008), 373–410.
- [2] R. F. Bailey and B. Stevens, Hamilton decompositions of complete k -uniform hypergraphs, *Discrete Math.* **310** (2010), 3088–3095.
- [3] Zs. Baranyai, On the factorization of the complete uniform hypergraph, in: Infinite and finite sets, *Colloq. Math. Soc. János Bolyai* **10**, North-Holland, Amsterdam, 1975, 91–108.
- [4] J.-C. Bermond, A. Germa, and D. Sotteau, Hypergraph-designs, *Ars Combinatoria* **3** (1977), 47–66.
- [5] D. Bryant, S. Herke, B. Maenhaut, and W. Wannasit, Decompositions of complete 3-uniform hypergraphs into small 3-uniform hypergraphs, *Australas. J. Combin.* **60** (2014), 227–254.
- [6] D. E. Bryant and T. A. McCourt, Existence results for G -designs, <http://wiki.smp.uq.edu.au/G-designs/>
- [7] R. C. Bunge, S. I. El-Zanati, J. Jetton, M. Juarez, A. J. Netz, D. Roberts, and P. Ward, On loose 5-cycle decompositions, packings, and coverings of complete lambda-fold 3-uniform hypergraphs, *Ars Combinatoria*, to appear.
- [8] C. J. Colbourn and R. Matheron, Steiner systems, in *The CRC Handbook of Combinatorial Designs*, 2nd edition, (Eds. C. J. Colbourn and J. H. Dinitz), CRC Press, Boca Raton (2007), 102–110.
- [9] S. Glock, D. Kühn, A. Lo, and D. Osthus, The existence of designs via iterative absorption, arXiv:1611.06827v2, (2017), 63 pages.
- [10] S. Glock, D. Kühn, A. Lo, and D. Osthus, Hypergraph F -designs for arbitrary F , arXiv:1706.01800, (2017), 72 pages.
- [11] H. Hanani, On quadruple systems, *Canad. J. Math.*, **12** (1960), 145–157.

- [12] H. Hanani, Decomposition of hypergraphs into octahedra, Second International Conference on Combinatorial Mathematics (New York, 1978), pp. 260–264, *Ann. New York Acad. Sci.*, 319, New York Acad. Sci., New York, 1979.
- [13] H. Jordon and G. Newkirk, 4-cycle decompositions of complete 3-uniform hypergraphs, *Australas. J. Combin.* **71** (2018), 312–323.
- [14] P. Keevash, The existence of designs, arXiv:1401.3665v2, (2018), 39 pages.
- [15] J. Kuhl and M. W. Schroeder, Hamilton cycle decompositions of k -uniform k -partite hypergraphs, *Australas. J. Combin.* **56** (2013), 23–37.
- [16] D. Kühn and D. Osthus, Decompositions of complete uniform hypergraphs into Hamilton Berge cycles, *J. Combin. Theory Ser. A* **126** (2014), 128–135.
- [17] M. Meszka and A. Rosa, Decomposing complete 3-uniform hypergraphs into Hamiltonian cycles, *Australas. J. Combin.* **45** (2009), 291–302.
- [18] M. W. Schroeder, On Hamilton cycle decompositions of r -uniform r -partite hypergraphs, *Discrete Math.* **315** (2014), 1–8.
- [19] R. M. Wilson, Decompositions of Complete Graphs into Subgraphs Isomorphic to a Given Graph, in “Proc. Fifth British Combinatorial Conference” (C. St. J. A. Nash-Williams and J. Sheehan, Eds.), pp. 647–659, *Congressus Numerantium XV*, 1975.